Rend. Mat. Appl. (7).
Volume 41 (2020), 105 – 116

RENDICONTI DI MATEMATICA
E DELLE SUE APPLICAZIONI

On the some subclasses of bi-univalent functions related to the Faber polynomial expansions and the Fibonacci numbers

Şahsene Altınkaya* **and Sibel Yalçın**

Abstract. *In this investigation, by using the Tremblay fractional derivative operator, we introduce the new class* $\mathfrak{I}_{\Sigma,\gamma}^{\mu,\rho}\left(\widetilde{\mathfrak{p}}\right)$ *of bi-univalent functions based on the rule of subordination. Moreover, we use the Faber polynomial expansions and Fibonacci numbers to derive bounds for the general coefficient* $|a_n|$ *of the bi-univalent function class.*

1. Introduction, Definitions and Notations

Let $\mathbb{C}$ be the complex plane and $\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$ be the open unit disc in $\mathbb{C}$. Further, let $\mathcal{A}$ represent the class of functions analytic in $\mathbb{U}$, satisfying the condition

$$f(0) = f'(0) - 1 = 0\,.$$

Then each function f in $\mathcal{A}$ has the following Taylor series expansion

$$f(z) = z + a_2 z^2 + a_3 z^3 + \cdots = z + \sum_{n=2}^{\infty} a_n z^n. \tag{1.1}$$

Let $\mathcal{S}$ represent the subclass of $\mathcal{A}$ consisting of functions univalent in $\mathbb{U}$. With a view to reminding the rule of subordination for analytic functions, let the functions f, g be analytic in $\mathbb{U}$. A function f is *subordinate* to g, indicated as $f \prec g$, if there exists a Schwarz function

$$\varpi(z) = \sum_{n=1}^{\infty} \mathfrak{c}_n z^n \quad (\varpi(0) = 0,\ |\varpi(z)| < 1)\,,$$

analytic in $\mathbb{U}$ such that

$$f(z) = g(\varpi(z)) \qquad (z \in \mathbb{U})\,.$$

For the Schwarz function $\varpi(z)$ we know that $|\mathfrak{c}_n| < 1$ (see [9]).

2020 Mathematics Subject Classification: Primary 30C45, 33D15.
Keywords: Bi-univalent functions, subordination, Faber polynomials, Fibonacci numbers, Tremblay fractional derivative operator.
© The Author(s) 2020. This article is an open access publication.
*Corresponding author.

According to the *Koebe-One Quarter Theorem*, every univalent function $f \in \mathcal{A}$ has an inverse f^{-1} satisfying $f^{-1}(f(z)) = z \quad (z \in \mathbb{U})$ and $f\left(f^{-1}(w)\right) = w$ $\left(|w| < r_0(f);\ \ r_0(f) \geq \frac{1}{4}\right)$, where

$$\begin{aligned} g(w) = f^{-1}(w) = w \ &- a_2 w^2 + \left(2a_2^2 - a_3\right) w^3 \\ &- \left(5a_2^3 - 5a_2 a_3 + a_4\right) w^4 + \cdots . \end{aligned} \tag{1.2}$$

A function $f \in \mathcal{A}$ is said to be bi-univalent in $\mathbb{U}$ if both f and f^{-1} are univalent in $\mathbb{U}$. Let Σ denote the class of bi-univalent functions in $\mathbb{U}$ given by (1.1). For a brief historical account and for several notable investigation of functions in the class Σ, see the pioneering work on this subject by Srivastava et al. [19] (see also [6, 7, 14, 15]). The interest on estimates for the first two coefficients $|a_2|$, $|a_3|$ of the bi-univalent functions keep on by many researchers (see, for example, [4, 12, 13, 20, 22, 23, 24]). However, in the literature, there are only a few works (by making use of the Faber polynomial expansions) determining the general coefficient bounds $|a_n|$ for bi-univalent functions ([5, 10, 11, 16, 21]). The coefficient estimate problem for each of $|a_n|$ ($n \in \mathbb{N}\backslash\{1,2\}$; $\mathbb{N} = \{1,2,3,\ldots\}$) is still an open problem.

Now, we recall to a notion of q-operators that play a major role in Geometric Function Theory. The application of the q-calculus in the context of Geometric Function Theory was actually provided and the basic (or q-) hypergeometric functions were first used in Geometric Function Theory in a book chapter by Srivastava [17]. For the convenience, we provide some basic notation details of q-calculus which are used in this paper.

Definition 1.1. (See [18]) For a function f (analytic in a simply-connected region of $\mathbb{C}$), the fractional derivative of order ρ is stated by

$$D_z^{\rho} f(z) = \frac{1}{\Gamma(1-\rho)} \frac{d}{dz} \int_0^z \frac{f(\xi)}{(z-\xi)^{\rho}} d\xi \quad (0 \leq \rho < 1)$$

and the fractional integral of order ρ is stated by

$$I_z^{\rho} f(z) = \frac{1}{\Gamma(\rho)} \int_0^z f(\xi)(z-\xi)^{\rho-1} d\xi \quad (\rho > 0).$$

Definition 1.2. (See [16]) The Tremblay fractional derivative operator of the function f is defined as

$$I_z^{\mu,\rho} f(z) = \frac{\Gamma(\rho)}{\Gamma(\mu)} z^{1-\rho} D_z^{\mu-\rho} z^{\mu-1} f(z) \tag{1.3}$$

$$(0 < \mu \leq 1,\ 0 < \rho \leq 1,\ \mu \geq \rho d,\ 0 < \mu - \rho < 1).$$

From (1.3), we deduce that

$$I_z^{\mu,\rho} f(z) = \frac{\mu}{\rho} z + \sum_{n=2}^{\infty} \frac{\Gamma(\rho)\Gamma(n+\mu)}{\Gamma(\mu)\Gamma(n+\rho)} a_n z^n.$$

In this paper, we study the new class $\mathfrak{I}_{\Sigma,\gamma}^{\mu,\rho}\left(\widetilde{\mathfrak{p}}\right)$ of bi-univalent functions established by using the Tremblay fractional derivative operator. Further, we use the Faber polynomial expansions and Fibonacci numbers to derive bounds for the general coefficient $|a_n|$ of the bi-univalent function class.

2. Preliminaries

By utilizing the Faber polynomial expansions for functions $f \in \mathcal{A}$ of the form (1.1), the coefficients of its inverse map $g = f^{-1}$ may be stated by (see [2, 3])

$$g(w) = f^{-1}(w) = w + \sum_{n=2}^{\infty} \frac{1}{n} K_{n-1}^{-n}(a_2, a_3, \ldots) w^n,$$

where

$$\begin{aligned} K_{n-1}^{-n} = {} & \frac{(-n)!}{(-2n+1)!\,(n-1)!} a_2^{n-1} + \frac{(-n)!}{[2(-n+1)]!\,(n-3)!} a_2^{n-3} a_3 \\ & + \frac{(-n)!}{(-2n+3)!\,(n-4)!} a_2^{n-4} a_4 \\ & + \frac{(-n)!}{[2(-n+2)]!\,(n-5)!} a_2^{n-5} \left[a_5 + (-n+2)\, a_3^2\right] \\ & + \frac{(-n)!}{(-2n+5)!\,(n-6)!} a_2^{n-6} \left[a_6 + (-2n+5)\, a_3 a_4\right] \\ & + \sum_{j \geq 7} a_2^{n-j} V_j, \end{aligned}$$

such that V_j $(7 \leq j \leq n)$ is a homogeneous polynomial in the variables $a_2, a_3, \ldots, a_n$. In the following, the first three terms of K_{n-1}^{-n} are stated by

$$\begin{aligned} \frac{1}{2} K_1^{-2} &= -a_2, \\ \frac{1}{3} K_2^{-3} &= 2a_2^2 - a_3, \\ \frac{1}{4} K_3^{-4} &= -\left(5a_2^3 - 5a_2 a_3 + a_4\right). \end{aligned}$$

In general, the expansion of K_n^p $(p \in \mathbb{Z} = \{0, \pm 1, \pm 2, \ldots\})$ is stated by

$$K_n^p = p a_n + \frac{p(p-1)}{2} \mathcal{G}_n^2 + \frac{p!}{(p-3)!3!} \mathcal{G}_n^3 + \cdots + \frac{p!}{(p-n)!n!} \mathcal{G}_n^n,$$

where $\mathcal{G}_n^p = \mathcal{G}_n^p(a_1, a_2, \ldots)$ and by [1],

$$\mathcal{G}_n^m(a_1, a_2, \ldots, a_n) = \sum_{n=1}^{\infty} \frac{m!\,(a_1)^{\delta_1} \cdots (a_n)^{\delta_n}}{\delta_1! \cdots \delta_n!},$$

while $a_1 = 1$, the sum is taken over all nonnegative integers $\delta_1, \ldots, \delta_n$ satisfying

$$\delta_1 + \delta_2 + \cdots + \delta_n = m,$$
$$\delta_1 + 2\delta_2 + \cdots + n\delta_n = n.$$

The first and the last polynomials are

$$\mathcal{G}_n^1 = a_n \qquad \mathcal{G}_n^n = a_1^n.$$

For two analytic functions $\mathfrak{u}(z)$, $\mathfrak{v}(w)$ $(\mathfrak{u}(0) = \mathfrak{v}(0) = 0,\ |\mathfrak{u}(z)| < 1,\ |\mathfrak{v}(w)| < 1)$, suppose that

$$\mathfrak{u}(z) = \textstyle\sum_{n=1}^{\infty} t_n z^n \quad (|z| < 1,\ z \in \mathbb{U}),$$

$$\mathfrak{v}(w) = \textstyle\sum_{n=1}^{\infty} s_n w^n \quad (|w| < 1,\ w \in \mathbb{U}).$$

It is well known that

$$|t_1| \leq 1, \quad |t_2| \leq 1 - |t_1|^2, \quad |s_1| \leq 1, \quad |s_2| \leq 1 - |s_1|^2. \tag{2.1}$$

Definition 2.1. A function $f \in \Sigma$ is said to be in the class

$$\mathfrak{I}_{\Sigma,\gamma}^{\mu,\rho}\left(\widetilde{\mathfrak{p}}\right) \quad (\gamma \geq 1,\ 0 < \mu \leq 1,\ 0 < \rho \leq 1,\ z, w \in \mathbb{U})$$

if the following subordination relationships are satisfied:

$$\left[\frac{(1-\gamma)\rho}{\mu}\frac{I_z^{\mu,\rho} f(z)}{z} + \frac{\rho\gamma}{\mu}\left(I_z^{\mu,\rho} f(z)\right)'\right] \prec \widetilde{\mathfrak{p}}(z) = \frac{1+\tau^2 z^2}{1-\tau z-\tau^2 z^2}$$

and

$$\left[\frac{(1-\gamma)\rho}{\mu}\frac{I_z^{\mu,\rho} g(w)}{w} + \frac{\rho\gamma}{\mu}\left(I_z^{\mu,\rho} g(w)\right)'\right] \prec \widetilde{\mathfrak{p}}(w) = \frac{1+\tau^2 w^2}{1-\tau w-\tau^2 w^2},$$

where the function g is given by (1.2) and $\tau = \frac{1-\sqrt{5}}{2} \approx -0.618$.

Remark 2.2. The function $\widetilde{\mathfrak{p}}(z)$ is not univalent in $\mathbb{U}$, but it is univalent in the disc $|z| < \frac{3-\sqrt{5}}{2} \approx 0.38$. For example, $\widetilde{\mathfrak{p}}(0) = \widetilde{\mathfrak{p}}\left(-\frac{1}{2\tau}\right)$ and $\widetilde{\mathfrak{p}}\left(e^{\pm i \arccos(1/4)}\right) = \frac{\sqrt{5}}{5}$. Also, it can be written as

$$\frac{1}{|\tau|} = \frac{|\tau|}{1-|\tau|}$$

which indicates that the number $|\tau|$ divides $[0, 1]$ such that it fulfills the golden section (see for details Dziok et al. [8]).

Additionally, Dziok et al. [8] indicate a useful connection between the function $\widetilde{\mathfrak{p}}(z)$ and the Fibonacci numbers. Let $\{\Lambda_n\}$ be the sequence of Fibonacci numbers

$$\Lambda_0 = 0, \quad \Lambda_1 = 1, \quad \Lambda_{n+2} = \Lambda_n + \Lambda_{n+1} \quad (n \in \mathbb{N}_0 = \{0, 1, 2, \ldots\}),$$

then

$$\Lambda_n = \frac{(1-\tau)^n - \tau^n}{\sqrt{5}}, \qquad \tau = \frac{1-\sqrt{5}}{2}.$$

If we set

$$\widetilde{\mathfrak{p}}(z) = 1 + \sum_{n=1}^{\infty} \widetilde{\mathfrak{p}}_n z^n = 1 + (\Lambda_0 + \Lambda_2)\tau z + (\Lambda_1 + \Lambda_3)\tau^2 z^2$$

$$+ \sum_{n=3}^{\infty} (\Lambda_{n-3} + \Lambda_{n-2} + \Lambda_{n-1} + \Lambda_n)\tau^n z^n,$$

then the coefficients $\widetilde{\mathfrak{p}}_n$ satisfy

$$\widetilde{\mathfrak{p}}_n = \begin{cases} \tau & (n = 1) \\ 3\tau^2 & (n = 2) \\ \tau\widetilde{\mathfrak{p}}_{n-1} + \tau^2\widetilde{\mathfrak{p}}_{n-2} & (n = 3, 4, \ldots) \end{cases} . \tag{2.2}$$

Specializing the parameters γ, μ and ρ, we state the following definitions.

Definition 2.3. For $\mu = \rho = 1$, a function $f \in \Sigma$ is said to be in the class $\mathfrak{I}_{\Sigma,\gamma}\left(\widetilde{\mathfrak{p}}\right)$ if it satisfies the following conditions respectively:

$$\left[(1-\gamma)\frac{f(z)}{z} + \gamma f'(z)\right] \prec \widetilde{\mathfrak{p}}(z)$$

and

$$\left[(1-\gamma)\frac{g(w)}{w} + \gamma g'(w)\right] \prec \widetilde{\mathfrak{p}}(w),$$

where $g = f^{-1}$.

Definition 2.4. For $\gamma = \mu = \rho = 1$, a function $f \in \Sigma$ is said to be in the class $\mathfrak{I}_{\Sigma}\left(\widetilde{\mathfrak{p}}\right)$ if it satisfies the following conditions respectively:

$$f'(z) \prec \widetilde{\mathfrak{p}}(z)$$

and

$$g'(w) \prec \widetilde{\mathfrak{p}}(w),$$

where $g = f^{-1}$.

3. Main Result and its consequences

Theorem 3.1. *For $\gamma \geq 1,\ 0 < \mu \leq 1$ and $0 < \rho \leq 1$, let the function f given by (1.1) be in the function class $\mathfrak{I}_{\Sigma,\gamma}^{\mu,\rho}\left(\widetilde{\mathfrak{p}}\right)$. If $a_m = 0\ \ (2 \leq m \leq n-1)$, then*

$$|a_n| \leq \frac{|\tau|\,\Gamma(\mu+1)\Gamma(n+\rho)}{\Gamma(\rho+1)\Gamma(n+\mu)\,[(n-1)\gamma+1]} \quad (n \geq 3).$$

Proof. By the definition of subordination yields

$$\left[\frac{(1-\gamma)\rho}{\mu}\frac{I_z^{\mu,\rho}f(z)}{z} + \frac{\rho\gamma}{\mu}\left(I_z^{\mu,\rho}f(z)\right)'\right] = \widetilde{\mathfrak{p}}(\mathfrak{u}(z)) \tag{3.1}$$

and

$$\left[\frac{(1-\gamma)\rho}{\mu}\frac{I_z^{\mu,\rho}g(w)}{w} + \frac{\rho\gamma}{\mu}\left(I_z^{\mu,\rho}g(w)\right)'\right] = \widetilde{\mathfrak{p}}(\mathfrak{v}(w)). \tag{3.2}$$

Using Faber polynomial expansions, we have

$$\frac{(1-\gamma)\rho}{\mu}\frac{I_z^{\mu,\rho}f(z)}{z}+\frac{\rho\gamma}{\mu}\left(I_z^{\mu,\rho}f(z)\right)' = 1+\sum_{n=2}^{\infty}\frac{\Gamma(\rho+1)\Gamma(n+\mu)}{\Gamma(\mu+1)\Gamma(n+\rho)}\left[(n-1)\gamma+1\right]a_n z^{n-1}$$

and for its inverse map $g = f^{-1}$, it is seen that

$$\begin{aligned}
&\frac{(1-\gamma)\rho}{\mu}\frac{I_z^{\mu,\rho}g(w)}{w} + \frac{\rho\gamma}{\mu}\left(I_z^{\mu,\rho}g(w)\right)' \\
&\quad = 1 + \sum_{n=2}^{\infty}\frac{\Gamma(\rho+1)\Gamma(n+\mu)}{\Gamma(\mu+1)\Gamma(n+\rho)}\left[(n-1)\gamma+1\right]\frac{1}{n}K_{n-1}^{-n}\left(a_2, a_3, \ldots\right)w^{n-1} \\
&\quad = 1 + \sum_{n=2}^{\infty}\frac{\Gamma(\rho+1)\Gamma(n+\mu)}{\Gamma(\mu+1)\Gamma(n+\rho)}\left[(n-1)\gamma+1\right]b_n w^{n-1}.
\end{aligned}$$

Next, the equations (3.1) and (3.2) lead to

$$\begin{aligned}
\widetilde{\mathfrak{p}}\left(\mathfrak{u}\left(z\right)\right) &= 1 + \widetilde{\mathfrak{p}}_1\mathfrak{u}(z) + \widetilde{\mathfrak{p}}_2(\mathfrak{u}(z))^2 z^2 + \cdots \\
&= 1 + \widetilde{\mathfrak{p}}_1 t_1 z + \left(\widetilde{\mathfrak{p}}_1 t_2 + \widetilde{\mathfrak{p}}_2 t_1^2\right) z^2 + \cdots \\
&= 1 + \sum_{n=1}^{\infty}\sum_{k=1}^{n}\widetilde{\mathfrak{p}}_k \mathcal{G}_n^k\left(t_1, t_2, \ldots, t_n\right) z^n,
\end{aligned}$$

and

$$\widetilde{\mathfrak{p}}\left(\mathfrak{v}\left(w\right)\right) = 1 + \widetilde{\mathfrak{p}}_1\mathfrak{v}(w) + \widetilde{\mathfrak{p}}_2(\mathfrak{v}(w))^2 w^2 + \cdots$$

$$= 1 + \widetilde{\mathfrak{p}}_1 s_1 w + \left(\widetilde{\mathfrak{p}}_1 s_2 + \widetilde{\mathfrak{p}}_2 s_1^2\right) w^2 + \cdots$$

$$= 1 + \sum_{n=1}^{\infty}\sum_{k=1}^{n}\widetilde{\mathfrak{p}}_k \mathcal{G}_n^k\left(s_1, s_2, \ldots, s_n\right) w^n.$$

Comparing the corresponding coefficients of (3.1) and (3.2) yields

$$\frac{\Gamma(\rho+1)\Gamma(n+\mu)}{\Gamma(\mu+1)\Gamma(n+\rho)}\left[(n-1)\gamma+1\right] a_n = \widetilde{\mathfrak{p}}_1 t_{n-1},$$

$$\frac{\Gamma(\rho+1)\Gamma(n+\mu)}{\Gamma(\mu+1)\Gamma(n+\rho)}\left[(n-1)\gamma+1\right] b_n = \widetilde{\mathfrak{p}}_1 s_{n-1}.$$

For $a_m = 0 \ \ (2 \leq m \leq n-1)$, we get $b_n = -a_n$ and so

$$\frac{\Gamma(\rho+1)\Gamma(n+\mu)}{\Gamma(\mu+1)\Gamma(n+\rho)}\left[(n-1)\gamma+1\right] a_n = \widetilde{\mathfrak{p}}_1 t_{n-1} \tag{3.3}$$

and

$$-\frac{\Gamma(\rho+1)\Gamma(n+\mu)}{\Gamma(\mu+1)\Gamma(n+\rho)}\left[(n-1)\gamma+1\right] a_n = \widetilde{\mathfrak{p}}_1 s_{n-1}. \tag{3.4}$$

Now taking the absolute values of either of the above two equations and from (2.1), we obtain

$$|a_n| \leq \frac{|\tau|\,\Gamma(\mu+1)\Gamma(n+\rho)}{\Gamma(\rho+1)\Gamma(n+\mu)\left[(n-1)\gamma+1\right]}.$$

□

Corollary 3.2. *For $\gamma \geq 1$, suppose that $f \in \mathfrak{I}_{\Sigma,\gamma}\left(\widetilde{\mathfrak{p}}\right)$. If $a_m = 0 \ \ (2 \leq m \leq n-1)$, then*

$$|a_n| \leq \frac{|\tau|}{\left[(n-1)\gamma+1\right]} \quad (n \geq 3).$$

Corollary 3.3. *Suppose that $f \in \mathfrak{I}_{\Sigma}\left(\widetilde{\mathfrak{p}}\right)$. If $a_m = 0 \ \ (2 \leq m \leq n-1)$, then*

$$|a_n| \leq \frac{|\tau|}{n} \quad (n \geq 3).$$

Theorem 3.4. *Let $f \in \mathfrak{I}_{\Sigma,\gamma}^{\mu,\rho}\left(\widetilde{\mathfrak{p}}\right)$. Then*

$$|a_2| \leq \min\left\{\frac{|\tau|}{\sqrt{\left|\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)} - \frac{3(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}\right||\tau| + \frac{(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}}}, \right.$$

$$\left. |\tau|\sqrt{\frac{3(\rho+1)(\rho+2)}{(\mu+1)(\mu+2)(2\gamma+1)}}\right\}$$

and

$$|a_3| \leq \min\left\{\frac{3\tau^2(\rho+1)(\rho+2)}{(\mu+1)(\mu+2)(2\gamma+1)},\right.$$
$$\left.\frac{|\tau|}{\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}}\left[1+\frac{\left[\frac{(\mu+1)(\mu+2)(2\gamma+1)|\tau|}{(\rho+1)(\rho+2)}-\frac{(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}\right]}{\left|\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}-\frac{3(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}\right||\tau|+\frac{(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}}\right]\right\}.$$

Proof. Substituting n by 2 and 3 in (3.3) and (3.4), respectively, we find that

$$\frac{(\mu+1)}{(\rho+1)}(\gamma+1)a_2 = \widetilde{\mathfrak{p}}_1 t_1, \tag{3.5}$$

$$\frac{(\mu+1)(\mu+2)}{(\rho+1)(\rho+2)}(2\gamma+1)a_3 = \widetilde{\mathfrak{p}}_1 t_2 + \widetilde{\mathfrak{p}}_2 t_1^2, \tag{3.6}$$

$$-\frac{(\mu+1)}{(\rho+1)}(\gamma+1)a_2 = \widetilde{\mathfrak{p}}_1 s_1, \tag{3.7}$$

$$\frac{(\mu+1)(\mu+2)}{(\rho+1)(\rho+2)}(2\gamma+1)(2a_2^2-a_3) = \widetilde{\mathfrak{p}}_1 s_2 + \widetilde{\mathfrak{p}}_2 s_1^2. \tag{3.8}$$

Obviously, we obtain

$$t_1 = -s_1. \tag{3.9}$$

If we add the equation (3.8) to (3.6) and use (3.9), we get

$$\frac{2(\mu+1)(\mu+2)}{(\rho+1)(\rho+2)}(2\gamma+1)a_2^2 = \widetilde{\mathfrak{p}}_1\left(t_2+s_2\right) + 2\widetilde{\mathfrak{p}}_2 t_1^2. \tag{3.10}$$

Using the value of t_1^2 from (3.5), we get

$$\left[\frac{2(\mu+1)(\mu+2)}{(\rho+1)(\rho+2)}(2\gamma+1)\widetilde{\mathfrak{p}}_1^2 - \frac{2(\mu+1)^2}{(\rho+1)^2}(\gamma+1)^2\widetilde{\mathfrak{p}}_2\right]a_2^2 = \widetilde{\mathfrak{p}}_1^3\left(t_2+s_2\right). \tag{3.11}$$

Combining (3.11) and (2.1), we obtain

$$\begin{aligned}2\left|\frac{(\mu+1)(\mu+2)}{(\rho+1)(\rho+2)}(2\gamma+1)\widetilde{\mathfrak{p}}_1^2 - \frac{(\mu+1)^2}{(\rho+1)^2}(\gamma+1)^2\widetilde{\mathfrak{p}}_2\right||a_2|^2 &\leq \left|\widetilde{\mathfrak{p}}_1\right|^3\left(|t_2|+|s_2|\right)\\ &\leq 2\left|\widetilde{\mathfrak{p}}_1\right|^3\left(1-|t_1|^2\right)\\ &= 2\left|\widetilde{\mathfrak{p}}_1\right|^3 - 2\left|\widetilde{\mathfrak{p}}_1\right|^3|t_1|^2.\end{aligned}$$

It follows from (3.5) that

$$|a_2| \leq \frac{|\tau|}{\sqrt{\left|\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}-\frac{3(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}\right||\tau|+\frac{(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}}}. \tag{3.12}$$

Additionally, by (2.1) and (3.10)

$$\begin{aligned}\frac{2(\mu+1)(\mu+2)}{(\rho+1)(\rho+2)}(2\gamma+1)\left|a_2\right|^2 &\leq \left|\widetilde{\mathfrak{p}}_1\right|(|t_2|+|s_2|)+2\left|\widetilde{\mathfrak{p}}_2\right|\left|t_1\right|^2\\ &\leq 2\left|\widetilde{\mathfrak{p}}_1\right|\left(1-\left|t_1\right|^2\right)+2\left|\widetilde{\mathfrak{p}}_2\right|\left|t_1\right|^2\\ &= 2\left|\widetilde{\mathfrak{p}}_1\right|+2\left|t_1\right|^2\left(\left|\widetilde{\mathfrak{p}}_2\right|-\left|\widetilde{\mathfrak{p}}_1\right|\right).\end{aligned}$$

Since $\left|\widetilde{\mathfrak{p}}_2\right| > \left|\widetilde{\mathfrak{p}}_1\right|$, we get

$$|a_2| \leq |\tau|\sqrt{\frac{3(\rho+1)(\rho+2)}{(\mu+1)(\mu+2)(2\gamma+1)}}.$$

Next, in order to derive the bounds on $|a_3|$, by subtracting (3.8) from (3.6), we may obtain

$$\frac{2(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}a_3 = \frac{2(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}a_2^2+\widetilde{\mathfrak{p}}_1\left(t_2-s_2\right). \quad (3.13)$$

Evidently, from (3.10), we state that

$$\begin{aligned}a_3 &= \frac{\widetilde{\mathfrak{p}}_1\left(t_2+s_2\right)+2\widetilde{\mathfrak{p}}_2 t_1^2}{\frac{2(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}}+\frac{\widetilde{\mathfrak{p}}_1\left(t_2-s_2\right)}{\frac{2(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}}\\ &= \frac{\widetilde{\mathfrak{p}}_1 t_2+\widetilde{\mathfrak{p}}_2 t_1^2}{\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}}\end{aligned}$$

and consequently

$$\begin{aligned}|a_3| &\leq \frac{\left|\widetilde{\mathfrak{p}}_1\right|\left|t_2\right|+\left|\widetilde{\mathfrak{p}}_2\right|\left|t_1\right|^2}{\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}}\\ &\leq \frac{\left|\widetilde{\mathfrak{p}}_1\right|\left(1-\left|t_1\right|^2\right)+\left|\widetilde{\mathfrak{p}}_2\right|\left|t_1\right|^2}{\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}}\\ &= \frac{\left|\widetilde{\mathfrak{p}}_1\right|+\left|t_1\right|^2\left(\left|\widetilde{\mathfrak{p}}_2\right|-\left|\widetilde{\mathfrak{p}}_1\right|\right)}{\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}}.\end{aligned}$$

Since $\left|\widetilde{\mathfrak{p}}_2\right| > \left|\widetilde{\mathfrak{p}}_1\right|$, we must write

$$|a_3| \leq \frac{3\tau^2(\rho+1)(\rho+2)}{(\mu+1)(\mu+2)(2\gamma+1)}.$$

On the other hand, by (2.1) and (3.13), we have

$$\frac{2(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}\left|a_3\right| \leq \frac{2(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}\left|a_2\right|^2+\left|\widetilde{\mathfrak{p}}_1\right|\left(\left|t_2\right|+\left|s_2\right|\right)$$

$$\leq \frac{2(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}\left|a_2\right|^2+2\left|\widetilde{\mathfrak{p}}_1\right|\left(1-\left|t_1\right|^2\right).$$

Then, with the help of (3.5), we have

$$\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}\left|a_3\right|$$

$$\leq \left[\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}-\frac{(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2\left|\widetilde{\mathfrak{p}}_1\right|}\right]\left|a_2\right|^2+\left|\widetilde{\mathfrak{p}}_1\right|.$$

By considering (3.12), we deduce that

$$\left|a_3\right| \leq \frac{|\tau|}{\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}}\left\{1+\frac{\left[\frac{(\mu+1)(\mu+2)(2\gamma+1)|\tau|}{(\rho+1)(\rho+2)}-\frac{(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}\right]}{\left|\frac{(\mu+1)(\mu+2)(2\gamma+1)}{(\rho+1)(\rho+2)}-\frac{3(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}\right||\tau|+\frac{(\mu+1)^2(\gamma+1)^2}{(\rho+1)^2}}\right\}.$$

□

Corollary 3.5. *Let $f \in \mathfrak{I}_{\Sigma,\gamma}\left(\widetilde{\mathfrak{p}}\right)$ $(\gamma \geq 1)$. Then*

$$\left|a_2\right| \leq \min\left\{\frac{|\tau|}{\sqrt{\left(3\gamma^2+4\gamma+2\right)|\tau|+(\gamma+1)^2}}, |\tau|\sqrt{\frac{3}{2\gamma+1}}\right\}$$

and

$$\left|a_3\right| \leq \min\left\{\frac{3\tau^2}{2\gamma+1}, \frac{|\tau|}{2\gamma+1}\left[1+\frac{(2\gamma+1)|\tau|-(\gamma+1)^2}{\left(3\gamma^2+4\gamma+2\right)|\tau|+(\gamma+1)^2}\right]\right\}.$$

Corollary 3.6. *Let $f \in \mathfrak{I}_{\Sigma}\left(\widetilde{\mathfrak{p}}\right)$. Then*

$$\left|a_2\right| \leq \frac{|\tau|}{\sqrt{9|\tau|+4}}$$

and

$$\left|a_3\right| \leq \frac{4|\tau|^2}{9|\tau|+4}.$$

Acknowledgements. The first author is supported by the Scientific and Technological Research Council of Turkey (TUBITAK 1002-Short Term R&D Funding Program) Project Number: 118F543.

References

[1] Airault, H.: Symmetric sums associated to the factorization of Grunsky coefficients. in Conference, Groups and Symmetries, Montreal, Canada, (2007)

[2] Airault, H., Bouali, H.: Differential calculus on the Faber polynomials. Bull. Sci. Math. **130**, 179-222 (2006)

[3] Airault, H., Ren, J.: An algebra of differential operators and generating functions on the set of univalent functions. Bull. Sci. Math. **126**, 343-367 (2002)

[4] Altınkaya, Ş., Yalçın, S.: Estimate for initial MacLaurin of general subclasses of bi-univalent functions of complex order involving subordination. Honam Math. J. **40**, 391-400 (2018)

[5] Altınkaya, Ş., Yalçın, S.: Faber polynomial coefficient estimates for bi-univalent functions of complex order based on subordinate conditions involving of the Jackson (p, q)-derivative operator. Miskolc Math. Notes **18**, 555-572 (2017)

[6] Brannan, D. A., Clunie, J. G.: Aspects of contemporary complex analysis. Proceedings of the NATO Advanced Study Institute Held at University of Durham, New York: Academic Press, (1979)

[7] Brannan, D. A., Taha, T. S.: On some classes of bi-univalent functions. Stud. Univ. Babeş-Bolyai Math. **31**, 70-77 (1986)

[8] Dziok, J., Raina, R. K., Sokół, J.: On α-convex functions related to shell-like functions connected with Fibonacci numbers. Appl. Math. Comput. **218**, 996–1002 (2011)

[9] Duren, P.L.: Univalent Functions, Grundlehren der Mathematischen Wissenschaften, Springer, New York, USA, **259**, (1983)

[10] Deniz, E., Jahangiri, J. M., Hamidi, S. G., Kına, S. S.: Faber polynomial coefficients for generalized bi-subordinate functions of complex order. J. Math. Inequal. **12**, 645–653 (2018)

[11] Hamidi, S. G., Jahangiri, J. M.: Faber polynomial coefficients of bi-subordinate functions. C. R. Acad. Sci. Paris, Ser. I **354**, 365-370 (2016)

[12] Hayami, T., Owa, S.: Coefficient bounds for bi-univalent functions. Pan Amer. Math. J. **22**, 15-26 (2012)

[13] Özlem Güney, H., Murugusundaramoorthy, G., Sokol, J.: Subclasses of bi-univalent functions related to shell-like curves connected with Fibonacci numbers. Acta Univ. Sapientiae Mathematica **10**, 70-84 (2018)

[14] Lewin, M.: On a coefficient problem for bi-univalent functions. Proc. Amer. Math. Soc. **18**, 63-68 (1967)

[15] Netanyahu, E.: The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $|z| < 1$. Arch. Ration. Mech. Anal. **32**, 100-112 (1969)

[16] Srivastava H. M., Sümer Eker, S., Hamidi, S. G., Jahangiri, J. M.: Faber polynomial coefficient estimates for bi-univalent functions defined by the Tremblay fractional derivative operator. Bull. Iranian Math. Soc. **44**, 149–157 (2018)

[17] Srivastava H. M.: Univalent functions, fractional calculus, and associated generalized hypergeometric functions, in Univalent Functions, Fractional Calculus, and Their Applications. (H. M. Srivastava and S. Owa, Editors), Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, (1989)

[18] Srivastava H. M., Owa, S.: Univalent Functions. Fractional Calculus, and Their Applications. Ellis Horwood Ltd. Publ., Chichester, (1989)

[19] Srivastava H. M., Mishra, A. K., Gochhayat, P.: Certain subclasses of analytic and bi-univalent functions. Appl. Math. Lett. **23**, 1188-1192 (2010)

[20] Srivastava H. M., Sakar, F. M., Güney, H. Ö.: Some general coefficient estimates for a new class of analytic and bi-univalent functions defined by a linear combination. Filomat **32**, 1313–1322 (2018)

[21] Srivastava H. M., Khan, S., Ahmad, Q.-Z., Khan, N., Hussain, S.:The Faber polynomial expansion method and its application to the general coefficient problem for some subclasses of bi-univalent functions associated with a certain q-integral operator. Stud. Univ. Babeş-Bolyai Math. **63**, 419-436 (2018)

[22] Srivastava H. M., Gaboury, S., Ghanim, F.: Coefficient estimates for a general subclass of analytic and bi-univalent functions of the Ma–Minda type. Rev. Real Acad. Cienc. Exactas Fis. Natur. Ser. A Mat. **112**, 1157-1168 (2018)

[23] Srivastava H. M., Wanas, A. K.: Initial Maclaurin coefficient bounds for new subclasses of analytic and m-fold symmetric bi-univalent functions defined by a linear combination. Kyungpook Math. J. **59**, 493-503 (2019)
[24] Srivastava H. M., Altınkaya Ş., Yalçın S.: Certain subclasses of Bi-univalent functions associated with the Horadam polynomials. Iran. J. Sci. Technol. Trans. A Sci. **43**, 1873-1879 (2019)

Received: 30 September 2019/Accepted: 8 January 2020/Published online: 9 January 2020

Şahsene Altınkaya
Department of Mathematics, Bursa Uludag University, Bursa 16059, Turkey.
sahsenealtinkaya@gmail.com

Sibel Yalçın
Department of Mathematics, Bursa Uludag University, Bursa 16059, Turkey.
syalcin@uludag.edu.tr

Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.