arXiv:1404.0905v1 [math.CA] 3 Apr 2014

ON NEW GENERAL INTEGRAL INEQUALITIES FOR s-CONVEX FUNCTIONS

İMDAT İŞCAN▼, ERHAN SET♣, AND M. EMIN ÖZDEMIR■

Abstract. In this paper, the authors establish some new estimates for the remainder term of the midpoint, trapezoid, and Simpson formula using functions whose derivatives in absolute value at certain power are s-convex. Some applications to special means of real numbers are provided as well.

1. Introduction

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$. The following inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2} \tag{1.1}$$

holds. This double inequality is known in the literature as Hermite-Hadamard integral inequality for convex functions. See ([2],[4],[6]-[12],[15]) for the results of the generalization, improvement and extention of the famous integral inequality (1.1).

In 1978, Breckner introduced s-convex functions as a generalization of convex functions as follows [3]:

Definition 1. *Let $s \in (0,1]$ be a fixed real number. A function $f : [0,\infty) \to [0,\infty)$ is said to be $s-$convex (in the second sense),or that f belongs to the class K_s^2, if*

$$f(\alpha x + (1-\alpha)y) \leq \alpha^s f(x) + (1-\alpha)^s f(y)$$

for all $x, y \in [0,\infty)$ and $\alpha \in [0,1]$.

Of course, s-convexity means just convexity when $s = 1$. For other recent results concerning s-convex functions see [1]-[19].

The following inequality is well known in the literature as Simpson's inequality:

Let $f : [a,b] \to \mathbb{R}$ be a four times continuously differentiable mapping on (a,b) and $\left\|f^{(4)}\right\|_\infty = \sup_{x\in(a,b)} \left|f^{(4)}(x)\right| < \infty$. Then the following inequality holds:

$$\left|\frac{1}{3}\left[\frac{f(a)+f(b)}{2} + 2f\left(\frac{a+b}{2}\right)\right] - \frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{1}{2880}\left\|f^{(4)}\right\|_\infty (b-a)^4.$$

2000 *Mathematics Subject Classification.* 26A51, 26D15.

Key words and phrases. convex function, s-convex function, Simpson's inequality, Hermite-Hadamard's inequality.

In recent years many authors have studied error estimations for Simpson's inequality. For refinements, counterparts, generalizations of the Simpson's inequality and new Simpson's type inequalities, see [1, 6, 7, 8, 13, 14, 19].

In [4], Dragomir and Fitzpatrick proved a variant of Hermite–Hadamard inequality which holds for the s-convex functions.

Theorem 1. *Suppose that $f:[0,\infty)\to[0,\infty)$ is an s-convex function in the second sense, where $s\in(0,1]$ and let $a,b\in[0,\infty)$, $a<b$. If $f\in L[a,b]$, then the following inequalities hold*

$$2^{s-1}f\left(\frac{a+b}{2}\right)\leq\frac{1}{b-a}\int_a^b f(x)dx\leq\frac{f(a)+f(b)}{s+1} \tag{1.2}$$

the constant $k=\frac{1}{s+1}$ is the best possible in the second inequality in (1.2). The above inequalities are sharp.

In [6], Iscan obtained a new generalization of some integral inequalities for differentiable convex mapping which are connected Simpson and Hadamard type inequalities, and he used the following lemma to prove this.

Lemma 1. *Let $f:I\subseteq\mathbb{R}\to\mathbb{R}$ be a differentiable mapping on I° such that $f'\in L[a,b]$, where $a,b\in I$ with $a<b$ and $\alpha,\lambda\in[0,1]$. Then the following equality holds:*

$$\begin{aligned}&\lambda\left(\alpha f(a)+(1-\alpha)f(b)\right)+(1-\lambda)f(\alpha a+(1-\alpha)b)-\frac{1}{b-a}\int_a^b f(x)dx\\ =\;&(b-a)\left[\int_0^{1-\alpha}(t-\alpha\lambda)f'(tb+(1-t)a)\,dt\right.\\ &\left.+\int_{1-\alpha}^{1}(t-1+\lambda(1-\alpha))f'(tb+(1-t)a)\,dt\right].\end{aligned}$$

The main inequality in [6], pointed out, is as follows.

Theorem 2. *Let $f:I\subseteq\mathbb{R}\to\mathbb{R}$ be a differentiable mapping on I° such that $f'\in L[a,b]$, where $a,b\in I^\circ$ with $a<b$ and $\alpha,\lambda\in[0,1]$. If $|f'|^q$ is convex on $[a,b]$, $q\geq1$, then the following inequality holds:*

$$\left|\lambda\left(\alpha f(a)+(1-\alpha)f(b)\right)+(1-\lambda)f(\alpha a+(1-\alpha)b)-\frac{1}{b-a}\int_a^b f(x)dx\right|$$

(1.3)

$$\leq \begin{cases} (b-a)\left\{\gamma_2^{1-\frac{1}{q}}\left(\mu_1\left|f'(b)\right|^q+\mu_2\left|f'(a)\right|^q\right)^{\frac{1}{q}}\right. & \\ \left.+\upsilon_2^{1-\frac{1}{q}}\left(\eta_3\left|f'(b)\right|^q+\eta_4\left|f'(a)\right|^q\right)^{\frac{1}{q}}\right\}, & \alpha\lambda\leq 1-\alpha\leq 1-\lambda(1-\alpha) \\ (b-a)\left\{\gamma_2^{1-\frac{1}{q}}\left(\mu_1\left|f'(b)\right|^q+\mu_2\left|f'(a)\right|^q\right)^{\frac{1}{q}}\right. & \\ \left.+\upsilon_1^{1-\frac{1}{q}}\left(\eta_1\left|f'(b)\right|^q+\eta_2\left|f'(a)\right|^q\right)^{\frac{1}{q}}\right\}, & \alpha\lambda\leq 1-\lambda(1-\alpha)\leq 1-\alpha \\ (b-a)\left\{\gamma_1^{1-\frac{1}{q}}\left(\mu_3\left|f'(b)\right|^q+\mu_4\left|f'(a)\right|^q\right)^{\frac{1}{q}}\right. & \\ \left.+\upsilon_2^{1-\frac{1}{q}}\left(\eta_3\left|f'(b)\right|^q+\eta_4\left|f'(a)\right|^q\right)^{\frac{1}{q}}\right\}, & 1-\alpha\leq\alpha\lambda\leq 1-\lambda(1-\alpha) \end{cases}$$

where

$$\gamma_1=(1-\alpha)\left[\alpha\lambda-\frac{(1-\alpha)}{2}\right],\ \gamma_2=(\alpha\lambda)^2-\gamma_1\ ,$$

$$\begin{aligned}\upsilon_1 &= \frac{1-(1-\alpha)^2}{2}-\alpha\left[1-\lambda(1-\alpha)\right],\\ \upsilon_2 &= \frac{1+(1-\alpha)^2}{2}-(\lambda+1)(1-\alpha)\left[1-\lambda(1-\alpha)\right],\end{aligned}$$

$$\begin{aligned}\mu_1 &= \frac{(\alpha\lambda)^3+(1-\alpha)^3}{3}-\alpha\lambda\frac{(1-\alpha)^2}{2},\\ \mu_2 &= \frac{1+\alpha^3+(1-\alpha\lambda)^3}{3}-\frac{(1-\alpha\lambda)}{2}\left(1+\alpha^2\right),\\ \mu_3 &= \alpha\lambda\frac{(1-\alpha)^2}{2}-\frac{(1-\alpha)^3}{3},\\ \mu_4 &= \frac{(\alpha\lambda-1)\left(1-\alpha^2\right)}{2}+\frac{1-\alpha^3}{3},\end{aligned}$$

$$\begin{aligned}\eta_1 &= \frac{1-(1-\alpha)^3}{3}-\frac{[1-\lambda(1-\alpha)]}{2}\alpha(2-\alpha),\\ \eta_2 &= \frac{\lambda(1-\alpha)\alpha^2}{2}-\frac{\alpha^3}{3},\end{aligned}$$

$$\begin{aligned}\eta_3 &= \frac{[1-\lambda(1-\alpha)]^3}{3}-\frac{[1-\lambda(1-\alpha)]}{2}\left(1+(1-\alpha)^2\right)+\frac{1+(1-\alpha)^3}{3},\\ \eta_4 &= \frac{[\lambda(1-\alpha)]^3}{3}-\frac{\lambda(1-\alpha)\alpha^2}{2}+\frac{\alpha^3}{3}.\end{aligned}$$

In [2] Alomari et al. obtained the following inequalities of the left-hand side of Hermite-Hadamard's inequality for s-convex mappings.

Theorem 3. *Let $f:I\subseteq[0,\infty)\to\mathbb{R}$ be a differentiable mapping on I°, such that $f'\in L[a,b]$, where $a,b\in I$ with $a<b$. If $|f'|^q$, $q\geq 1$, is s-convex on $[a,b]$, for*

some fixed $s \in (0,1]$, then the following inequality holds:

$$\begin{aligned}&\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right| \\ \leq\ & \frac{b-a}{8}\left(\frac{2}{(s+1)(s+2)}\right)^{\frac{1}{q}}\left[\left\{\left(2^{1-s}+1\right)|f'(b)|^q+2^{1-s}|f'(a)|^q\right\}^{\frac{1}{q}}\right. \\ &\left.+\left\{\left(2^{1-s}+1\right)|f'(a)|^q+2^{1-s}|f'(b)|^q\right\}^{\frac{1}{q}}\right].\end{aligned} \tag{1.4}$$

Theorem 4. *Let $f: I \subseteq [0,\infty) \to \mathbb{R}$ be a differentiable mapping on I°, such that $f' \in L[a,b]$, where $a,b \in I$ with $a<b$. If $|f'|^{\frac{p}{p-1}}$, $p>1$, is s-convex on $[a,b]$, for some fixed $s \in (0,1]$, then the following inequality holds:*

$$\begin{aligned}\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right| \ \leq\ & \left(\frac{b-a}{4}\right)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{s+1}\right)^{\frac{2}{q}} \\ & \times\left[\left(\left(2^{1-s}+s+1\right)|f'(a)|^q+2^{1-s}|f'(b)|^q\right)^{\frac{1}{q}}\right. \\ & \left.+\left(\left(2^{1-s}+s+1\right)|f'(b)|^q+2^{1-s}|f'(a)|^q\right)^{\frac{1}{q}}\right],\end{aligned} \tag{1.5}$$

where p is the conjugate of q, $q=p/(p-1)$.

In [14], Sarikaya et al. obtained a new upper bound for the right-hand side of Simpson's inequality for $s-$convex mapping as follows:

Theorem 5. *Let $f: I \subseteq [0,\infty) \to \mathbb{R}$ be a differentiable mapping on I°, such that $f' \in L[a,b]$, where $a,b \in I^\circ$ with $a<b$. If $|f'|^q$, is s-convex on $[a,b]$, for some fixed $s \in (0,1]$ and $q>1$, then the following inequality holds:*

$$\begin{aligned}&\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{12}\left(\frac{1+2^{p+1}}{3(p+1)}\right)^{\frac{1}{p}} \\ &\times\left\{\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(a)|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(b)|^q}{s+1}\right)^{\frac{1}{q}}\right\},\end{aligned} \tag{1.6}$$

where $\frac{1}{p}+\frac{1}{q}=1$.

In [10], Kirmaci et al. proved the following trapezoid inequality:

Theorem 6. *Let $f: I \subseteq [0,\infty) \to \mathbb{R}$ be a differentiable mapping on I°, such that $f' \in L[a,b]$, where $a,b \in I^\circ$, $a<b$. If $|f'|^q$, is s-convex on $[a,b]$, for some fixed $s \in (0,1)$ and $q>1$, then*

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{2}\left(\frac{q-1}{2(2q-1)}\right)^{\frac{q-1}{q}}\left(\frac{1}{s+1}\right)^{\frac{1}{q}} \\ &\times\left\{\left(\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(a)|^q\right)^{\frac{1}{q}}+\left(\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(b)|^q\right)^{\frac{1}{q}}\right\}.\end{aligned} \tag{1.7}$$

2. Main results

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable function on I°, the interior of I, throughout this section we will take

$$\begin{aligned} & I_f(\lambda,\alpha,a,b) \\ = \;& \lambda(\alpha f(a)+(1-\alpha)f(b))+(1-\lambda)f(\alpha a+(1-\alpha)b)-\frac{1}{b-a}\int_a^b f(x)dx \end{aligned}$$

where $a,b \in I^\circ$ with $a<b$ and $\alpha,\lambda \in [0,1]$.

Theorem 7. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a,b]$, where $a,b \in I^\circ$ with $a<b$ and $\alpha,\lambda \in [0,1]$. If $|f'|^q$ is s-convex on $[a,b]$, for some fixed $s \in (0,1]$ and $q \geq 1$, then*

(i) for $\alpha\lambda \leq 1-\alpha \leq 1-\lambda(1-\alpha)$ we have

$$\begin{aligned} |I_f(\lambda,\alpha,a,b)| \leq\;& (b-a)\left[\gamma_2^{1-\frac{1}{q}}(\alpha,\lambda)\left(c_1(\alpha,\lambda,s)|f'(b)|^q+c_2(\alpha,\lambda,s)|f'(a)|^q\right)^{\frac{1}{q}}\right. \\ &\left.+\gamma_2^{1-\frac{1}{q}}(1-\alpha,\lambda)\left(c_2(1-\alpha,\lambda,s)|f'(b)|^q+c_1(1-\alpha,\lambda,s)|f'(a)|^q\right)^{\frac{1}{q}}\right], \end{aligned}$$

(ii) for $\alpha\lambda \leq 1-\lambda(1-\alpha) \leq 1-\alpha$ we have

$$\begin{aligned} |I_f(\lambda,\alpha,a,b)| \leq\;& (b-a)\left[\gamma_2^{1-\frac{1}{q}}(\alpha,\lambda)\left(c_1(\alpha,\lambda,s)|f'(b)|^q+c_2(\alpha,\lambda,s)|f'(a)|^q\right)^{\frac{1}{q}}\right. \\ &\left.+\gamma_1^{1-\frac{1}{q}}(1-\alpha,\lambda)\left(c_4(1-\alpha,\lambda,s)|f'(b)|^q+c_3(1-\alpha,\lambda,s)|f'(a)|^q\right)^{\frac{1}{q}}\right], \end{aligned}$$

(iii) for $1-\alpha \leq \alpha\lambda \leq 1-\lambda(1-\alpha)$ we have

$$\begin{aligned} |I_f(\lambda,\alpha,a,b)| \leq\;& (b-a)\left[\gamma_1^{1-\frac{1}{q}}(\alpha,\lambda)\left(c_3(\alpha,\lambda,s)|f'(b)|^q+c_4(\alpha,\lambda,s)|f'(a)|^q\right)^{\frac{1}{q}}\right. \\ &\left.+\gamma_2^{1-\frac{1}{q}}(1-\alpha,\lambda)\left(c_2(1-\alpha,\lambda,s)|f'(b)|^q+c_1(1-\alpha,\lambda,s)|f'(a)|^q\right)^{\frac{1}{q}}\right] \end{aligned}$$

where

$$\begin{aligned} \gamma_1(\alpha,\lambda) &= (1-\alpha)\left[\alpha\lambda-\frac{(1-\alpha)}{2}\right], \\ \gamma_2(\alpha,\lambda) &= (\alpha\lambda)^2-\gamma_1(\alpha,\lambda), \end{aligned}$$

$$\begin{aligned} c_1(\alpha,\lambda,s) &= (\alpha\lambda)^{s+2}\frac{2}{(s+1)(s+2)}-(\alpha\lambda)\frac{(1-\alpha)^{s+1}}{s+1}+\frac{(1-\alpha)^{s+2}}{s+2}, \\ c_2(\alpha,\lambda,s) &= (1-\alpha\lambda)^{s+2}\frac{2}{(s+1)(s+2)}-\frac{(1-\alpha\lambda)\left(1+\alpha^{s+1}\right)}{s+1}+\frac{1+\alpha^{s+2}}{s+2}, \\ c_3(\alpha,\lambda,s) &= (\alpha\lambda)\frac{(1-\alpha)^{s+1}}{s+1}-\frac{(1-\alpha)^{s+2}}{s+2}, \\ c_4(\alpha,\lambda,s) &= \frac{(\alpha\lambda-1)\left(1-\alpha^{s+1}\right)}{s+1}+\frac{1-\alpha^{s+2}}{s+2}. \end{aligned}$$

Proof. Suppose that $q \geq 1$. From Lemma 1 and using the well known power mean inequality, we have

$$\begin{aligned}
&|I_f(\lambda,\alpha,a,b)| \\
\leq\ & (b-a)\left[\int_0^{1-\alpha}|t-\alpha\lambda|\,|f'(tb+(1-t)a)|\,dt+\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|\,|f'(tb+(1-t)a)|\,dt\right] \\
\leq\ & (b-a)\left\{\left(\int_0^{1-\alpha}|t-\alpha\lambda|\,dt\right)^{1-\frac{1}{q}}\left(\int_0^{1-\alpha}|t-\alpha\lambda|\,|f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\right. \\
&\left.+\left(\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|\,dt\right)^{1-\frac{1}{q}}\left(\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|\,|f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\right\}
\end{aligned} \tag{2.1}$$

Consider

$$I_1=\int_0^{1-\alpha}|t-\alpha\lambda|\,|f'(tb+(1-t)a)|^q\,dt,\ \ I_2=\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|\,|f'(tb+(1-t)a)|^q\,dt$$

Since $|f'|^q$ is s-convex on $[a,b]$,

$$I_1\leq|f'(b)|^q\int_0^{1-\alpha}|t-\alpha\lambda|\,t^s dt+|f'(a)|^q\int_0^{1-\alpha}|t-\alpha\lambda|\,(1-t)^s dt. \tag{2.2}$$

Similarly

$$I_2\leq|f'(b)|^q\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|\,t^s dt+|f'(a)|^q\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|\,(1-t)^s dt. \tag{2.3}$$

Additionally, by simple computation

$$\int_0^{1-\alpha}|t-\alpha\lambda|\,dt=\begin{cases}\gamma_2(\alpha,\lambda), & \alpha\lambda\leq 1-\alpha\\ \gamma_1(\alpha,\lambda), & \alpha\lambda\geq 1-\alpha\end{cases}, \tag{2.4}$$

$$\gamma_1(\alpha,\lambda)=(1-\alpha)\left[\alpha\lambda-\frac{(1-\alpha)}{2}\right],\ \gamma_2(\alpha,\lambda)=(\alpha\lambda)^2-\gamma_1(\alpha,\lambda)\ ,$$

$$\begin{aligned}
\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|\,dt&=\int_0^{\alpha}|t-(1-\alpha)\lambda|\,dt\\
&=\begin{cases}\gamma_1(1-\alpha,\lambda), & 1-\lambda(1-\alpha)\leq 1-\alpha\\ \gamma_2(1-\alpha,\lambda), & 1-\lambda(1-\alpha)\geq 1-\alpha\end{cases},
\end{aligned}$$

$$\int_0^{1-\alpha}|t-\alpha\lambda|\,t^s dt=\begin{cases}c_1(\alpha,\lambda,s), & \alpha\lambda\leq 1-\alpha\\ c_3(\alpha,\lambda,s), & \alpha\lambda\geq 1-\alpha\end{cases}$$

$$\int_0^{1-\alpha} |t-\alpha\lambda|\,(1-t)^s dt = \begin{cases} c_2(\alpha,\lambda,s), & \alpha\lambda \le 1-\alpha \\ c_4(\alpha,\lambda,s), & \alpha\lambda \ge 1-\alpha \end{cases}$$

$$\int_{1-\alpha}^{1} |t-1+\lambda(1-\alpha)|\,t^s = \int_0^{\alpha} |t-(1-\alpha)\lambda|\,(1-t)^s dt$$

$$= \begin{cases} c_4(1-\alpha,\lambda,s), & 1-\lambda(1-\alpha) \le 1-\alpha \\ c_2(1-\alpha,\lambda,s), & 1-\lambda(1-\alpha) \ge 1-\alpha \end{cases}$$

$$\int_{1-\alpha}^{1} |t-1+\lambda(1-\alpha)|\,(1-t)^s dt = \int_0^{\alpha} |t-(1-\alpha)\lambda|\,t^s dt$$

$$= \begin{cases} c_3(1-\alpha,\lambda,s), & 1-\lambda(1-\alpha) \le 1-\alpha \\ c_1(1-\alpha,\lambda,s), & 1-\lambda(1-\alpha) \ge 1-\alpha \end{cases}. \tag{2.5}$$

Thus, using (2.2)-(2.5) in (2.1), we obtain desired results. This completes the proof. □

Corollary 1. *Under the assumptions of Theorem 7 with $q=1$,*

(i) if $\alpha\lambda \le 1-\alpha \le 1-\lambda(1-\alpha)$, then we have

$$\begin{aligned} |I_f(\lambda,\alpha,a,b)| &\le (b-a)\left[\left(c_1(\alpha,\lambda,s)+c_2(1-\alpha,\lambda,s)\right)|f'(b)| \right. \\ &\quad \left. + \left(c_2(\alpha,\lambda,s)+c_1(1-\alpha,\lambda,s)\right)|f'(a)|\right], \end{aligned}$$

(ii) if $\alpha\lambda \le 1-\lambda(1-\alpha) \le 1-\alpha$, then we have

$$\begin{aligned} |I_f(\lambda,\alpha,a,b)| &\le (b-a)\left[\left(c_1(\alpha,\lambda,s)+c_4(1-\alpha,\lambda,s)\right)|f'(b)| \right. \\ &\quad \left. + \left(c_2(\alpha,\lambda,s)+c_3(1-\alpha,\lambda,s)\right)|f'(a)|\right], \end{aligned}$$

(iii) if $1-\alpha \le \alpha\lambda \le 1-\lambda(1-\alpha)$, then we have

$$\begin{aligned} |I_f(\lambda,\alpha,a,b)| &\le (b-a)\left[\left(c_3(\alpha,\lambda,s)+c_2(1-\alpha,\lambda,s)\right)|f'(b)| \right. \\ &\quad \left. + \left(c_4(\alpha,\lambda,s)+c_1(1-\alpha,\lambda,s)\right)|f'(a)|\right] \end{aligned}$$

Remark 1. *In Theorem 7, if we take $s=1$, then we obtain the inequality (1.3).*

Remark 2. *In Theorem 7, if we take $\alpha=\frac{1}{2}$ and $\lambda=\frac{1}{3}$, then we have the following Simpson type inequality*

$$\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{b-a}\int_a^b f(x)dx\right| \le \frac{b-a}{2}\left(\frac{5}{36}\right)^{1-\frac{1}{q}} \tag{2.6}$$

$$\times\left\{\left(\frac{(2s+1)3^{s+1}+2}{3\times 6^{s+1}(s+1)(s+2)}|f'(b)|^q+\frac{2\times 5^{s+2}+(s-4)6^{s+1}-(2s+7)3^{s+1}}{3\times 6^{s+1}(s+1)(s+2)}|f'(a)|^q\right)^{\frac{1}{q}}\right.$$

$$\left.+\left(\frac{2\times 5^{s+2}+(s-4)6^{s+1}-(2s+7)3^{s+1}}{3.6^{s+1}(s+1)(s+2)}|f'(b)|^q+\frac{(2s+1)3^{s+1}+2}{3\times 6^{s+1}(s+1)(s+2)}|f'(a)|^q\right)^{\frac{1}{q}}\right\},$$

which is the same of the inequality in [14, Theorem 10] .

Remark 3. *In Theorem 7 , if we take $\alpha = \frac{1}{2}$ and $\lambda = 0$, then we have following midpoint inequality*

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a}\int_a^b f(x)dx \right| \leq \frac{b-a}{8}\left(\frac{2}{(s+1)(s+2)}\right)^{\frac{1}{q}} \times \left\{ \left(\frac{2^{1-s}(s+1)|f'(b)|^q}{2} + \frac{2^{1-s}\left(2^{s+2}-s-3\right)|f'(a)|^q}{2} \right)^{\frac{1}{q}} + \left(\frac{2^{1-s}(s+1)|f'(a)|^q}{2} + \frac{2^{1-s}\left(2^{s+2}-s-3\right)|f'(b)|^q}{2} \right)^{\frac{1}{q}} \right\}. \tag{2.7}$$

We note that the obtained midpoint inequality (2.7) is better than the inequality (1.4). Because $\frac{s+1}{2} \leq 1$ and $\frac{2^{s+2}-s-3}{2} \leq \frac{2^{1-s}+1}{2^{1-s}}$.

Remark 4. *In Theorem 7 , if we take $\alpha = \frac{1}{2}$, and $\lambda = 1$, then we get the following trapezoid inequality*

$$\left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)dx \right| \leq \frac{b-a}{8}\left(\frac{2^{1-s}}{(s+1)(s+2)}\right)^{\frac{1}{q}} \times \left\{ \left(|f'(b)|^q + |f'(a)|^q\left(2^{s+1}+1\right)\right)^{\frac{1}{q}} + \left(|f'(a)|^q + |f'(b)|^q\left(2^{s+1}+1\right)\right)^{\frac{1}{q}} \right\}$$

Using Lemma 1 we shall give another result for convex functions as follows.

Theorem 8. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a,b]$, where $a, b \in I^\circ$ with $a < b$ and $\alpha, \lambda \in [0,1]$. If $|f'|^q$ is s-convex on $[a,b]$, for some fixed $s \in (0,1]$ and $q > 1$, then*

$$|I_f(\lambda,\alpha,a,b)| \leq (b-a)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{s+1}\right)^{\frac{1}{q}} \times \begin{cases} \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)C_f^{1/q}(\alpha,q) + \varepsilon_1^{1/p}(1-\alpha,\lambda,p)D_f^{1/q}(\alpha,q)\right], & \alpha\lambda \leq 1-\alpha \leq 1-\lambda(1-\alpha) \\ \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)C_f^{1/q}(\alpha,q) + \varepsilon_2^{1/p}(1-\alpha,\lambda,p)D_f^{1/q}(\alpha,q)\right], & \alpha\lambda \leq 1-\lambda(1-\alpha) \leq 1-\alpha \\ \left[\varepsilon_2^{1/p}(\alpha,\lambda,p)C_f^{1/q}(\alpha,q) + \varepsilon_1^{1/p}(1-\alpha,\lambda,p)D_f^{1/q}(\alpha,q)\right], & 1-\alpha \leq \alpha\lambda \leq 1-\lambda(1-\alpha) \end{cases}, \tag{2.8}$$

where

$$\begin{aligned} C_f(\alpha,q) &= (1-\alpha)\left[|f'((1-\alpha)b+\alpha a)|^q + |f'(a)|^q\right], \\ D_f(\alpha,q) &= \alpha\left[|f'((1-\alpha)b+\alpha a)|^q + |f'(b)|^q\right], \end{aligned} \tag{2.9}$$

$$\begin{aligned} \varepsilon_1(\alpha,\lambda,p) &= (\alpha\lambda)^{p+1} + (1-\alpha-\alpha\lambda)^{p+1}, \\ \varepsilon_2(\alpha,\lambda,p) &= (\alpha\lambda)^{p+1} - (\alpha\lambda-1+\alpha)^{p+1}, \end{aligned} \tag{2.10}$$

and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From Lemma 1 and by Hölder's integral inequality, we have

$$\begin{aligned}
&|I_f(\lambda,\alpha,a,b)| \\
\leq\ & (b-a)\left[\int_0^{1-\alpha}|t-\alpha\lambda|\,|f'(tb+(1-t)a)|\,dt+\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|\,|f'(tb+(1-t)a)|\,dt\right] \\
\leq\ & (b-a)\left\{\left(\int_0^{1-\alpha}|t-\alpha\lambda|^p\,dt\right)^{\frac{1}{p}}\left(\int_0^{1-\alpha}|f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\right. \\
&\left.+\left(\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|^p\,dt\right)^{\frac{1}{p}}\left(\int_{1-\alpha}^{1}|f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\right\}. \qquad (2.11)
\end{aligned}$$

Since $|f'|^q$ is s-convex on $[a,b]$, for $\alpha\in[0,1)$ by the inequality (1.2), we get

$$\begin{aligned}
\int_0^{1-\alpha}|f'(tb+(1-t)a)|^q\,dt &= (1-\alpha)\left[\frac{1}{(1-\alpha)(b-a)}\int_a^{(1-\alpha)b+\alpha a}|f'(x)|^q\,dx\right] \\
&\leq (1-\alpha)\left[\frac{|f'((1-\alpha)b+\alpha a)|^q+|f'(a)|^q}{s+1}\right]. \qquad (2.12)
\end{aligned}$$

The inequality (2.12) also holds for $\alpha=1$. Similarly, for $\alpha\in(0,1]$ by the inequality (1.2), we have

$$\begin{aligned}
\int_{1-\alpha}^{1}|f'(tb+(1-t)a)|^q\,dt &= \alpha\left[\frac{1}{\alpha(b-a)}\int_{(1-\alpha)b+\alpha a}^{b}|f'(x)|^q\,dx\right] \\
&\leq \alpha\left[\frac{|f'((1-\alpha)b+\alpha a)|^q+|f'(b)|^q}{s+1}\right]. \qquad (2.13)
\end{aligned}$$

The inequality (2.13) also holds for $\alpha=0$. By simple computation

$$\int_0^{1-\alpha}|t-\alpha\lambda|^p\,dt=\begin{cases}\frac{(\alpha\lambda)^{p+1}+(1-\alpha-\alpha\lambda)^{p+1}}{p+1}, & \alpha\lambda\leq 1-\alpha\\ \frac{(\alpha\lambda)^{p+1}-(\alpha\lambda-1+\alpha)^{p+1}}{p+1}, & \alpha\lambda\geq 1-\alpha\end{cases}, \qquad (2.14)$$

and

$$\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|^p\,dt=\begin{cases}\frac{[\lambda(1-\alpha)]^{p+1}+[\alpha-\lambda(1-\alpha)]^{p+1}}{p+1}, & 1-\alpha\leq 1-\lambda(1-\alpha)\\ \frac{[\lambda(1-\alpha)]^{p+1}-[\lambda(1-\alpha)-\alpha]^{p+1}}{p+1}, & 1-\alpha\geq 1-\lambda(1-\alpha)\end{cases}, \qquad (2.15)$$

thus, using (2.12)-(2.15) in (2.11), we obtain the inequality (2.8). This completes the proof. □

Corollary 2. *Under the assumptions of Theorem 8 with $s=1$, we have*

$$|I_f(\lambda,\alpha,a,b)|\leq(b-a)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1}{q}}$$

$$\times \begin{cases} \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)C_f^{1/q}(\alpha,q)+\varepsilon_1^{1/p}(1-\alpha,\lambda,p)D_f^{1/q}(\alpha,q)\right], & \alpha\lambda \le 1-\alpha \le 1-\lambda(1-\alpha) \\ \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)C_f^{1/q}(\alpha,q)+\varepsilon_2^{1/p}(1-\alpha,\lambda,p)D_f^{1/q}(\alpha,q)\right], & \alpha\lambda \le 1-\lambda(1-\alpha) \le 1-\alpha \\ \left[\varepsilon_2^{1/p}(\alpha,\lambda,p)C_f^{1/q}(\alpha,q)+\varepsilon_1^{1/p}(1-\alpha,\lambda,p)D_f^{1/q}(\alpha,q)\right], & 1-\alpha \le \alpha\lambda \le 1-\lambda(1-\alpha) \end{cases},$$

where ε_1, ε_2, C_f and D_f are defined as in (2.9).

Remark 5. *In Theorem 8, if we take $\alpha=\frac{1}{2}$ and $\lambda=\frac{1}{3}$, then we have the following Simpson type inequality*

$$\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{b-a}\int_a^b f(x)dx\right| \tag{2.16}$$

$$\le \frac{b-a}{12}\left(\frac{1+2^{p+1}}{3(p+1)}\right)^{\frac{1}{p}}\left\{\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(a)|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(b)|^q}{s+1}\right)^{\frac{1}{q}}\right\},$$

which is the same of the inequality (1.6).

Remark 6. *In Theorem 8, if we take $\alpha=\frac{1}{2}$ and $\lambda=0$, then we have the following midpoint inequality*

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right|$$

$$\le \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left\{\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(a)|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(b)|^q}{s+1}\right)^{\frac{1}{q}}\right\}.$$

We note that by inequality

$$2^{s-1}\left|f'\left(\frac{a+b}{2}\right)\right|^q \le \frac{|f'(a)|^q+|f'(b)|^q}{s+1}$$

we have

$$\begin{aligned}\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right| &\le \left(\frac{b-a}{4}\right)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{s+1}\right)^{\frac{2}{q}} \\ &\quad\times\left[\left(\left(2^{1-s}+s+1\right)|f'(a)|^q+2^{1-s}|f'(b)|^q\right)^{\frac{1}{q}}\right. \\ &\quad\left.+\left(\left(2^{1-s}+s+1\right)|f'(b)|^q+2^{1-s}|f'(a)|^q\right)^{\frac{1}{q}}\right],\end{aligned}$$

which is the same of the inequality (1.5).

Remark 7. *In Theorem 8, if we take $\alpha=\frac{1}{2}$ and $\lambda=1$, then we have the following trapezoid inequality*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right| \le \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \tag{2.17}$$

$$\times\left\{\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(a)|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(b)|^q}{s+1}\right)^{\frac{1}{q}}\right\}.$$

We note that the obtained midpoint inequality (2.17) is better than the inequality (1.7).

Theorem 9. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a,b]$, where $a, b \in I^\circ$ with $a < b$ and $\alpha, \lambda \in [0,1]$. If $|f'|^q$ is s-concave on $[a,b]$, for some fixed $s \in (0,1]$ and $q > 1$, then the following inequality holds:*

$$|I_f(\lambda,\alpha,a,b)| \leq (b-a)\, 2^{\frac{s-1}{q}} \left(\frac{1}{p+1}\right)^{\frac{1}{p}} \tag{2.18}$$

$$\times \begin{cases} \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)E_f^{1/q}(\alpha,q) + \varepsilon_1^{1/p}(1-\alpha,\lambda,p)F_f^{1/q}(\alpha,q)\right], & \alpha\lambda \leq 1-\alpha \leq 1-\lambda(1-\alpha) \\ \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)E_f^{1/q}(\alpha,q) + \varepsilon_2^{1/p}(1-\alpha,\lambda,p)F_f^{1/q}(\alpha,q)\right], & \alpha\lambda \leq 1-\lambda(1-\alpha) \leq 1-\alpha \\ \left[\varepsilon_2^{1/p}(\alpha,\lambda,p)E_f^{1/q}(\alpha,q) + \varepsilon_1^{1/p}(1-\alpha,\lambda,p)F_f^{1/q}(\alpha,q)\right], & 1-\alpha \leq \alpha\lambda \leq 1-\lambda(1-\alpha) \end{cases},$$

where

$$E_f(\alpha,q) = (1-\alpha)\left|f'\left(\frac{(1-\alpha)\,b + (1+\alpha)\,a}{2}\right)\right|^q,\ F_f(\alpha,q) = \alpha\left|f'\left(\frac{(2-\alpha)\,b + \alpha a}{2}\right)\right|^q,$$

and $\varepsilon_1,\ \varepsilon_2$ are defined as in (2.9).

Proof. We proceed similarly as in the proof Theorem 8. Since $|f'|^q$ is $s-$concave on $[a,b]$, for $\alpha \in [0,1)$ by the inequality (1.2), we get

$$\begin{aligned} \int_0^{1-\alpha} |f'(tb+(1-t)a)|^q\, dt &= (1-\alpha)\left[\frac{1}{(1-\alpha)(b-a)} \int_a^{(1-\alpha)b+\alpha a} |f'(x)|^q\, dx\right] \\ &\leq 2^{s-1}(1-\alpha)\left|f'\left(\frac{(1-\alpha)\,b + (1+\alpha)\,a}{2}\right)\right|^q \end{aligned} \tag{2.19}$$

The inequality (2.19) also holds for $\alpha = 1$. Similarly, for $\alpha \in (0,1]$ by the inequality (1.2), we have

$$\begin{aligned} \int_{1-\alpha}^{1} |f'(tb+(1-t)a)|^q\, dt &= \alpha\left[\frac{1}{\alpha(b-a)} \int_{(1-\alpha)b+\alpha a}^{b} |f'(x)|^q\, dx\right] \\ &\leq 2^{s-1}\alpha\left|f'\left(\frac{(2-\alpha)\,b + \alpha a}{2}\right)\right|^q \end{aligned} \tag{2.20}$$

The inequality (2.20) also holds for $\alpha = 0$. Thus, using (2.14),(2.15),(2.19)and (2.20) in (2.11), we obtain the inequality (2.18). This completes the proof. □

Corollary 3. *Under the assumptions of Theorem 9 with $s = 1$, we have*

$$|I_f(\lambda,\alpha,a,b)| \leq (b-a)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}$$

$$\times \begin{cases} \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)E_f^{1/q}(\alpha,q) + \varepsilon_1^{1/p}(1-\alpha,\lambda,p)F_f^{1/q}(\alpha,q)\right], & \alpha\lambda \leq 1-\alpha \leq 1-\lambda(1-\alpha) \\ \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)E_f^{1/q}(\alpha,q) + \varepsilon_2^{1/p}(1-\alpha,\lambda,p)F_f^{1/q}(\alpha,q)\right], & \alpha\lambda \leq 1-\lambda(1-\alpha) \leq 1-\alpha \\ \left[\varepsilon_2^{1/p}(\alpha,\lambda,p)E_f^{1/q}(\alpha,q) + \varepsilon_1^{1/p}(1-\alpha,\lambda,p)F_f^{1/q}(\alpha,q)\right], & 1-\alpha \leq \alpha\lambda \leq 1-\lambda(1-\alpha) \end{cases},$$

where $\varepsilon_1,\ \varepsilon_2,\ E_f$ and F_f are defined as in Theorem 9.

Remark 8. *In Theorem 9, if we take* $\alpha = \frac{1}{2}$ *and* $\lambda = 1$*, then we have the following trapezoid inequality*

$$\begin{aligned} &\left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)dx \right| \\ \leq\ & \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \times \left(\frac{1}{2}\right)^{\frac{1-s}{q}} \left[\left| f'\left(\frac{3b+a}{4}\right)\right| + \left| f'\left(\frac{3a+b}{4}\right)\right| \right] \end{aligned}$$

which is the same of the inequality in [12, Theorem 8 (i)].

Remark 9. *In Theorem 9, if we take* $\alpha = \frac{1}{2}$ *and* $\lambda = 0$*, then we have the following midpoint inequality*

$$\begin{aligned} &\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a}\int_a^b f(x)dx \right| \\ \leq\ & \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \times \left(\frac{1}{2}\right)^{\frac{1-s}{q}} \left[\left| f'\left(\frac{3b+a}{4}\right)\right| + \left| f'\left(\frac{3a+b}{4}\right)\right| \right] \end{aligned}$$

which is the same of the inequality in [12, Theorem 8 (ii)].

Remark 10. *In Theorem 9, if we take* $\alpha = \frac{1}{2}$ *and* $\lambda = 1$*, then we have the following trapezoid inequality*

$$\begin{aligned} &\left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)dx \right| \\ \leq\ & \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \left[\left| f'\left(\frac{3b+a}{4}\right)\right| + \left| f'\left(\frac{3a+b}{4}\right)\right| \right] \end{aligned} \tag{2.21}$$

which is the same of the inequality in [10, Theorem 2].

Remark 11. *In Theorem 9, if we take* $\alpha = \frac{1}{2}$ *and* $\lambda = 0$*, then we have the following trapezoid inequality*

$$\begin{aligned} &\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a}\int_a^b f(x)dx \right| \\ \leq\ & \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \left[\left| f'\left(\frac{3b+a}{4}\right)\right| + \left| f'\left(\frac{3a+b}{4}\right)\right| \right] \end{aligned} \tag{2.22}$$

which is the same of the inequality in [2, Theorem 2.5].

Remark 12. *In Theorem 9, since* $|f'|^q$, $q > 1$*, is concave on* $[a,b]$*, using the power mean inequality, we have*

$$\begin{aligned} |f'(\lambda x + (1-\lambda)y)|^q &\geq \lambda |f'(x)|^q + (1-\lambda)|f'(y)|^q \\ &\geq (\lambda |f'(x)| + (1-\lambda)|f'(y)|)^q, \end{aligned}$$

$\forall x, y \in [a,b]$ *and* $\lambda \in [0,1]$*. Hence*

$$|f'(\lambda x + (1-\lambda)y)| \geq \lambda |f'(x)| + (1-\lambda)|f'(y)|$$

so $|f'|$ is also concave. Then by the inequality (1.1), we have

$$\left|f'\left(\frac{3b+a}{4}\right)\right|+\left|f'\left(\frac{3a+b}{4}\right)\right|\leq 2\left|f'\left(\frac{a+b}{2}\right)\right|. \tag{2.23}$$

Thus, using the inequality (2.23) in (2.21) and (2.22) we get

$$\begin{aligned}\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right| &\leq \frac{b-a}{2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left|f'\left(\frac{a+b}{2}\right)\right|,\\ \left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right| &\leq \frac{b-a}{2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left|f'\left(\frac{a+b}{2}\right)\right|.\end{aligned}$$

3. Some applications for special means

Let us recall the following special means of arbitrary real numbers a,b with $a\neq b$ and $\alpha\in[0,1]$:

(1) The weighted arithmetic mean

$$A_\alpha(a,b):=\alpha a+(1-\alpha)b,\ a,b\in\mathbb{R}.$$

(2) The unweighted arithmetic mean

$$A(a,b):=\frac{a+b}{2},\ a,b\in\mathbb{R}.$$

(3) Then $p-$Logarithmic mean

$$L_p(a,b):=\left(\frac{b^{p+1}-a^{p+1}}{(p+1)(b-a)}\right)^{\frac{1}{p}},\ p\in\mathbb{R}\backslash\{-1,0\},\ a,b>0.$$

From known Example 1 in [5], we may find that for any $s\in(0,1)$ and $\beta>0$, $f:[0,\infty)\to[0,\infty)$, $f(t)=\beta t^s$, $f\in K_s^2$.

Now, using the resuls of Section 2, some new inequalities are derived for the above means.

Proposition 1. *Let $a,b\in\mathbb{R}$ with $0<a<b$, $q\geq 1$ and $s\in\left(0,\frac{1}{q}\right)$. Then*

Theorem 10. *(i) for $\alpha\lambda\leq 1-\alpha\leq 1-\lambda(1-\alpha)$ we have*

$$\begin{aligned}&\left|\lambda A_\alpha\left(a^{s+1},b^{s+1}\right)+(1-\lambda)A_\alpha^{s+1}(a,b)-L_{s+1}^{s+1}(a,b)\right|\\ \leq\ &(b-a)(s+1)\left[\gamma_2^{1-\frac{1}{q}}(\alpha,\lambda)\left(c_1(\alpha,\lambda,s)b^{sq}+c_2(\alpha,\lambda,s)a^{sq}\right)^{\frac{1}{q}}\right.\\ &\left.+\gamma_2^{1-\frac{1}{q}}(1-\alpha,\lambda)\left(c_2(1-\alpha,\lambda,s)b^{sq}+c_1(1-\alpha,\lambda,s)a^{sq}\right)^{\frac{1}{q}}\right],\end{aligned}$$

(ii) for $\alpha\lambda\leq 1-\lambda(1-\alpha)\leq 1-\alpha$ we have

$$\begin{aligned}&\left|\lambda A_\alpha\left(a^{s+1},b^{s+1}\right)+(1-\lambda)A_\alpha^{s+1}(a,b)-L_{s+1}^{s+1}(a,b)\right|\\ \leq\ &(b-a)(s+1)\left[\gamma_2^{1-\frac{1}{q}}(\alpha,\lambda)\left(c_1(\alpha,\lambda,s)b^{sq}+c_2(\alpha,\lambda,s)a^{sq}\right)^{\frac{1}{q}}\right.\\ &\left.+\gamma_1^{1-\frac{1}{q}}(1-\alpha,\lambda)\left(c_4(1-\alpha,\lambda,s)b^{sq}+c_3(1-\alpha,\lambda,s)a^{sq}\right)^{\frac{1}{q}}\right],\end{aligned}$$

(iii) for $1-\alpha \leq \alpha\lambda \leq 1-\lambda(1-\alpha)$ *we have*

$$\begin{aligned} & \left|\lambda A_\alpha\left(a^{s+1},b^{s+1}\right)+(1-\lambda)A_\alpha^{s+1}(a,b)-L_{s+1}^{s+1}(a,b)\right| \\ \leq\ & (b-a)(s+1)\left[\gamma_1^{1-\frac{1}{q}}(\alpha,\lambda)\left(c_3(\alpha,\lambda,s)b^{sq}+c_4(\alpha,\lambda,s)a^{sq}\right)^{\frac{1}{q}}\right. \\ & \left.+\gamma_2^{1-\frac{1}{q}}(1-\alpha,\lambda)\left(c_2(1-\alpha,\lambda,s)b^{sq}+c_1(1-\alpha,\lambda,s)a^{sq}\right)^{\frac{1}{q}}\right] \end{aligned}$$

where $\gamma_1,\ \gamma_2,\ c_1,\ \ c_2,\ c_3,\ c_4$ *numbers are defined as in Theorem 7.*

Proof. The assertion follows from applied the inequalities in Theorem 7 to the function $f(t)=t^{s+1}$, $t\in[a,b]$ and $s\in\left(0,\frac{1}{q}\right)$, which implies that $f'(t)=(s+1)t^s,\ \ t\in[a,b]$ and $|f'(t)|^q=(s+1)^q t^{qs},\ \ t\in[a,b]$ is a s-convex function in the second sense since $qs\in(0,1)$ and $(s+1)^q>0$. □

Proposition 2. *Let* $a,b\in\mathbb{R}$ *with* $0<a<b,\ p,q>1,\ \frac{1}{p}+\frac{1}{q}=1$ *and* $s\in\left(0,\frac{1}{q}\right)$ *we have the following inequality:*

$$\left|\lambda A_\alpha\left(a^{s+1},b^{s+1}\right)+(1-\lambda)A_\alpha^{s+1}(a,b)-L_{s+1}^{s+1}(a,b)\right|\leq(b-a)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}(s+1)^{1-\frac{1}{q}}$$

$$\times\begin{cases}\left[\varepsilon_1^{1/p}(\alpha,\lambda,p)C_s^{1/q}(\alpha,q)+\varepsilon_1^{1/p}(1-\alpha,\lambda,p)D_s^{1/q}(\alpha,q)\right], & \alpha\lambda\leq 1-\alpha\leq 1-\lambda(1-\alpha)\\ \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)C_s^{1/q}(\alpha,q)+\varepsilon_2^{1/p}(1-\alpha,\lambda,p)D_s^{1/q}(\alpha,q)\right], & \alpha\lambda\leq 1-\lambda(1-\alpha)\leq 1-\alpha\\ \left[\varepsilon_2^{1/p}(\alpha,\lambda,p)C_s^{1/q}(\alpha,q)+\varepsilon_1^{1/p}(1-\alpha,\lambda,p)D_s^{1/q}(\alpha,q)\right], & 1-\alpha\leq\alpha\lambda\leq 1-\lambda(1-\alpha)\end{cases},$$

where

$$C_s(\alpha,q)=(1-\alpha)\left[A_\alpha^{sq}(a,b)+a^{sq}\right],\ D_s(\alpha,q)=\alpha\left[A_\alpha^{sq}(a,b)+b^{sq}\right],$$

and ε_1 *and* ε_2 *numbers are defined as in (2.10).*

Proof. The assertion follows from applied the inequality (2.8) to the function $f(t)=t^{s+1}$, $t\in[a,b]$ and $s\in\left(0,\frac{1}{q}\right)$, which implies that $f'(t)=(s+1)t^s,\ \ t\in[a,b]$ and $|f'(t)|^q=(s+1)^q t^{qs},\ \ t\in[a,b]$ is a s-convex function in the second sense since $qs\in(0,1)$ and $(s+1)^q>0$. □

References

[1] M. Alomari, M. Darus, S.S. Dragomir, New inequalities of Simpson's Type for s-convex functions with applications, RGMIA Res. Rep. Coll. **12**(4) (2009) Article 9. Online http://ajmaa.org/RGMIA/v12n4.php.

[2] M.W. Alomari, M. Darus, U.S. Kirmaci, Some inequalities of Hermite-Hadamard type for s-convex functions, Acta Math. Sci. **31B** (4) (2011), 1643-1652.

[3] W.W. Breckner, Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer funktionen in topologischen linearen Räumen, Publ. Inst. Math. **23** (1978), 13-20.

[4] S.S. Dragomir, S. Fitzpatrik, The Hadamard's inequality for s-convex functions in the second sense, Demonstr. Math.32(4), (1999), 687-696.

[5] H. Hudzik and L. Maligranda, Some remarks on s-convex functions, Aequationes Math. **48**(1) (1998), 100-111.

[6] I. Iscan, A new generalization of some integral inequalities and their applications, International Journal of Engineering and Applied sciences, **3**(3) (2013), 17-27.

[7] I. Iscan, Generalization of different type integral inequalities for s-convex functions via fractional integrals, Applicable Analysis, doi:10.1080/00036811.2013.851785. Available online at: http://dx.doi.org/10.1080/00036811.2013.851785

[8] I. Iscan, New estimates on generalization of some integral inequalities for s-convex functions and their applications, Int. J. Pure Appl. Math. **86**(4) (2013), 727-746.
[9] U.S. Kirmaci, Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, Appl. Math. Comput. **147** (2004), 137-146.
[10] U.S. Kirmaci, M.K. Bakula, M.E. Özdemir, J. Pečarić, Hadamard-type inequalities for s-convex functions, Appl. Math. Comput. **193** (2007), 26-35.
[11] Z. Liu, A note on Ostrowski type inequalities related to some s-convex functions in the second sense, Bull. Korean Math. Soc. **49**(4) (2012), 775-785.
[12] J. Park, Hermite-Hadamard-type inequalities for real α-star s-convex mappings, J. Appl. Math. & Informatics, **28(**5-6) (2010), 1507-1518.
[13] M.Z. Sarikaya, N. Aktan, On the generalization of some integral inequalities and their applications, Math. Comput. Modelling **54** (2011) 2175-2182.
[14] M.Z. Sarikaya, E. Set, M.E. Özdemir, On new inequalities of Simpson's type for s-convex functions, Comput. Math. Appl. **60** (2010) 2191-2199.
[15] Sarıkaya M.Z., Set E., Ögülmüş H., Some new inequalities of Hermite-Hadamard type for mappings whose derivatives are s-convex in the second sense, Int. J. of Modern Math. Sci., **8** (3) (2013), 212-218.
[16] E. Set, New inequalities of Ostrowski type for mappings whose derivatives are s-convex in the second sense via fractional integrals, Comput. Math. Appl., **63** (2012), 1147-1154.
[17] Set E., Özdemir M.E., Sarıkaya M.Z., New inequalities of Ostrowski's type for s-convex functions in the second sense with applications, Facta Universitatis Ser. Math. Inform., **27**(1) (2012), 67-82.
[18] Özdemir M.E., Yıldız Ç., Akdemir A.O., Set E., On some inequalities for s-convex functions and applications, Journal of Inequalities and Applications, 2013 (1):333, 11 pages (2013).
[19] Sarıkaya M.Z., Set E., Özdemir M.E., On new inequalities of Simpson's type for functions whose second derivatives absolute values are convex, Journal of Appl. Math., Statis. and Inform., **9** (1) (2013), 37-45.

▼Department of Mathematics, Faculty of Arts and Sciences, Giresun University, 28100, Giresun, Turkey.
E-mail address: imdat.iscan@giresun.edu.tr, imdati@yahoo.com

♣Department of Mathematics, Faculty of Arts and Sciences, Ordu University, 52200, Ordu, Turkey
E-mail address: erhanset@yahoo.com

■Atatürk University, K.K. Education Faculty, Department of Mathematics, 25240, Campus, Erzurum, Turkey
E-mail address: emos@atauni.edu.tr