FCMS
Fundamentals of Contemporary Mathematical Sciences
(**2021**) **2**(1) 52 – 69

New Integral Inequalities for Co-Ordinated Convex Functions

M. E. Özdemir [1], **A. O. Akdemir** [2*], **A. Ekinci** [3]
[1] Bursa Uludağ University, Faculty of Education, Department of Mathematics Education, Bursa, Türkiye, eminozdemir@uludag.edu.tr
[2] Ağrı İbrahim Çeçen University, Faculty of Arts and Sciences, Department of Mathematics Ağrı, Türkiye
[3] Bandırma Onyedi Eylül University, Bandırma Vocational School Department of International Trade, Bandırma, Türkiye, alperekinci@hotmail.com

Received: 19 December 2020 **Accepted:** 27 January 2021

Abstract: In this paper, we prove some new integral inequalities for co-ordinated convex functions by using a new lemma and fairly elementary analysis.

Keywords: Co-ordinated convex functions, Hadamard-type inequalities, Hölder inequality, power mean inequality.

1. Introduction

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a < b$. The following double inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}$$

is well known in the literature as Hadamard's inequality. Several papers have been written on convexity and inequalities (see [16–27] and references there in).

In [13], Dragomir defined convex functions on the co-ordinates as following:

Definition 1.1 *Let us consider the bidimensional interval* $\Delta = [a,b] \times [c,d]$ *in* $\mathbb{R}^2$ *with* $a < b$, $c < d$. *A function* $f : \Delta \to \mathbb{R}$ *will be called convex on the co-ordinates if the partial mappings* $f_y : [a,b] \to \mathbb{R}$, $f_y(u) = f(u,y)$ *and* $f_x : [c,d] \to \mathbb{R}$, $f_x(v) = f(x,v)$ *are convex where defined for all* $y \in [c,d]$ *and* $x \in [a,b]$. *Recall that the mapping* $f : \Delta \to \mathbb{R}$ *is convex on* Δ *if the following inequality holds,*

$$f(\lambda x + (1-\lambda)z, \lambda y + (1-\lambda)w) \leq \lambda f(x,y) + (1-\lambda)f(z,w)$$

*Correspondence: ahmetakdemir@agri.edu.tr
2020 *AMS Mathematics Subject Classification*: 26D15

This article is licensed under a Creative Commons Attribution 4.0 International License.
Also, it has been published considering the Research and Publication Ethics.

for all $(x,y),(z,w) \in \Delta$ *and* $\lambda \in [0,1]$.

In [13], Dragomir established the following inequalities of Hadamard's type for co-ordinated convex functions on a rectangle from the plane $\mathbb{R}^2$.

Theorem 1.2 *Suppose that* $f : \Delta = [a,b] \times [c,d] \to \mathbb{R}$ *is convex on the co-ordinates on* Δ. *Then one has the inequalities;*

$$\begin{aligned} & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \qquad (1) \\ \leq\ & \frac{1}{2}\left[\frac{1}{b-a}\int_a^b f\left(x, \frac{c+d}{2}\right)dx + \frac{1}{d-c}\int_c^d f\left(\frac{a+b}{2}, y\right)dy\right] \\ \leq\ & \frac{1}{(b-a)(d-c)}\int_a^b \int_c^d f(x,y)dxdy \\ \leq\ & \frac{1}{4}\left[\frac{1}{(b-a)}\int_a^b f(x,c)dx + \frac{1}{(b-a)}\int_a^b f(x,d)dx \right. \\ & \left. + \frac{1}{(d-c)}\int_c^d f(a,y)dy + \frac{1}{(d-c)}\int_c^d f(b,y)dy\right] \\ \leq\ & \frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}. \end{aligned}$$

The above inequalities are sharp.

In [1], Bakula and Pečarić established several Jensen type inequalities for co-ordinated convex functions and in [14], Hwang *et al.* gave a mapping F, discussed some properties of this mapping and proved some Hadamard-type inequalities for Lipschizian mapping in two variables. In [2], Özdemir *et al.* established new Hadamard-type inequalities for co-ordinated $m-$convex and $(\alpha,m)-$convex functions. On all of these, in [11], Sarıkaya *et al.* proved some Hadamard-type inequalities for co-ordinated convex functions as followings:

Theorem 1.3 *Let* $f : \Delta \subset \mathbb{R}^2 \to \mathbb{R}$ *be a partial differentiable mapping on* $\Delta := [a,b] \times [c,d]$ *in* $\mathbb{R}^2$ *with* $a < b$ *and* $c < d$. *If* $\left|\frac{\partial^2 f}{\partial t \partial s}\right|$ *is a convex function on the co-ordinates on* Δ, *then one has the*

inequalities:

$$\begin{aligned} &\left|\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right. \\ &\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy-A\right| \\ \leq\ & \frac{(b-a)(d-c)}{16}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|}{4}\right) \end{aligned} \tag{2}$$

where

$$A=\frac{1}{2}\left[\frac{1}{(b-a)}\int_a^b[f(x,c)+f(x,d)]\,dx+\frac{1}{(d-c)}\int_c^d[f(a,y)dy+f(b,y)]\,dy\right].$$

Theorem 1.4 *Let $f:\Delta\subset\mathbb{R}^2\to\mathbb{R}$ be a partial differentiable mapping on $\Delta:=[a,b]\times[c,d]$ in $\mathbb{R}^2$ with $a<b$ and $c<d$. If $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$, $q>1$, is a convex function on the co-ordinates on Δ, then one has the inequalities:*

$$\begin{aligned} &\left|\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right. \\ &\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy-A\right| \\ \leq\ & \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4}\right)^{\frac{1}{q}} \end{aligned} \tag{3}$$

where

$$A=\frac{1}{2}\left[\frac{1}{(b-a)}\int_a^b[f(x,c)+f(x,d)]\,dx+\frac{1}{(d-c)}\int_c^d[f(a,y)dy+f(b,y)]\,dy\right]$$

and $\frac{1}{p}+\frac{1}{q}=1$.

Theorem 1.5 *Let $f:\Delta\subset\mathbb{R}^2\to\mathbb{R}$ be a partial differentiable mapping on $\Delta:=[a,b]\times[c,d]$ in $\mathbb{R}^2$ with $a<b$ and $c<d$. If $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$, $q\geq 1$, is a convex function on the co-ordinates on Δ, then one*

has the inequalities:

$$\begin{aligned} &\left|\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right. \\ &\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy-A\right| \\ \leq\ &\frac{(b-a)(d-c)}{16}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4}\right)^{\frac{1}{q}} \end{aligned} \tag{4}$$

where

$$A=\frac{1}{2}\left[\frac{1}{(b-a)}\int_a^b\left[f(x,c)+f(x,d)\right]dx+\frac{1}{(d-c)}\int_c^d\left[f(a,y)dy+f(b,y)\right]dy\right].$$

In [15], Özdemir *et al.* proved following inequalities for co-ordinated convex functions.

Theorem 1.6 *Let $f:\Delta=[a,b]\times[c,d]\to\mathbb{R}$ be a partial differentiable mapping on $\Delta=[a,b]\times[c,d]$. If $\left|\frac{\partial^2 f}{\partial t\partial s}\right|$ is a convex function on the co-ordinates on Δ, then the following inequality holds;*

$$\begin{aligned} &\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right. \\ &-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx \\ &\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right| \\ \leq\ &\frac{(b-a)(d-c)}{64}\left[\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|\right]. \end{aligned} \tag{5}$$

Remark 1.7 *Suppose that all the assumptions of Theorem 5 are satisfied. If we choose $\frac{\partial^2 f}{\partial t\partial s}$ is bounded, i.e.,*

$$\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty=\sup_{(t,s)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right|<\infty,$$

we get

$$\begin{aligned} &\left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right)\right. \\ &-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2}, y\right)dy - \frac{1}{(b-a)}\int_a^b f\left(x, \frac{c+d}{2}\right)dx \\ &\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right| \\ \leq\ & \frac{(b-a)(d-c)}{16}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty . \end{aligned} \tag{6}$$

Theorem 1.8 *Let $f:\Delta=[a,b]\times[c,d]\to\mathbb{R}$ be a partial differentiable mapping on $\Delta=[a,b]\times[c,d]$. If $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$, $q>1$, is a convex function on the co-ordinates on Δ, then the following inequality holds;*

$$\begin{aligned} &\left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) + \frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right. \\ &\left.-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2}, y\right)dy - \frac{1}{(b-a)}\int_a^b f\left(x, \frac{c+d}{2}\right)dx\right| \\ \leq\ & \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}} \\ &\times\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4}\right)^{\frac{1}{q}} . \end{aligned} \tag{7}$$

Remark 1.9 *Suppose that all the assumptions of Theorem 6 are satisfied. If we choose $\frac{\partial^2 f}{\partial t\partial s}$ is bounded, i.e.,*

$$\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty = \sup_{(t,s)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right| < \infty,$$

we get

$$\begin{aligned} &\left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right)\right. \\ &-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2}, y\right)dy - \frac{1}{(b-a)}\int_a^b f\left(x, \frac{c+d}{2}\right)dx \\ &\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right| \\ \leq\ & \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty . \end{aligned} \tag{8}$$

Theorem 1.10 *Let $f : \Delta = [a,b] \times [c,d] \to \mathbb{R}$ be a partial differentiable mapping on $\Delta = [a,b] \times [c,d]$. If $\left|\frac{\partial^2 f}{\partial t \partial s}\right|^q$, $q \geq 1$, is a convex function on the co-ordinates on Δ, then the following inequality holds;*

$$\begin{aligned} &\left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x,y)\, dydx \right. \\ &\left. -\frac{1}{(d-c)} \int_c^d f\left(\frac{a+b}{2}, y\right) dy - \frac{1}{(b-a)} \int_a^b f\left(x, \frac{c+d}{2}\right) dx \right| \\ \leq\ & \frac{(b-a)(d-c)}{16} \\ &\times \left(\frac{\left|\frac{\partial^2 f}{\partial t \partial s}(a,c)\right|^q + \left|\frac{\partial^2 f}{\partial t \partial s}(b,c)\right|^q + \left|\frac{\partial^2 f}{\partial t \partial s}(a,d)\right|^q + \left|\frac{\partial^2 f}{\partial t \partial s}(b,d)\right|^q}{4} \right)^{\frac{1}{q}}. \end{aligned} \tag{9}$$

Recent results, generalizations, new methods and further properties of different kinds of co-ordinated convex functions can be found in [3–12].

The main purpose of this paper is to prove some new inequalities of Hadamard-type for co-ordinated convex functions on the co-ordinates by using a new Lemma. These estimations give new upper bounds.

2. Main Results

To prove our main results, we need following Lemma.

Lemma 2.1 *Let $f : \Delta = [a,b] \times [c,d] \to \mathbb{R}$ be a partial differentiable mapping on $\Delta = [a,b] \times [c,d]$. If $\frac{\partial^2 f}{\partial t \partial s} \in L(\Delta)$, then the following equality holds:*

$$\begin{aligned} \Lambda &= f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\ &-\frac{1}{(d-c)} \int_c^d f\left(\frac{a+b}{2}, y\right) dy - \frac{1}{(b-a)} \int_a^b f\left(x, \frac{c+d}{2}\right) dx \\ &+\frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x,y)\, dydx \end{aligned}$$

$$
\begin{aligned}
= \quad & \frac{(b-a)(d-c)}{16}\left[\int_0^1\int_0^1 ts\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)dsdt\right.\\
& +\int_0^1\int_0^1 (t-1)(s-1)\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right)dsdt\\
& +\int_0^1\int_0^1 t(s-1)\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, sd+(1-s)\frac{c+d}{2}\right)dsdt\\
& \left.+\int_0^1\int_0^1 s(t-1)\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)dsdt\right]
\end{aligned}
$$

for all $t,s\in[0,1]$.

Proof Let we start with the following statement

$$
I_1=\int_0^1\int_0^1 ts\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)dsdt.
$$

Integration by parts, we can write

$$
\begin{aligned}
& \int_0^1\int_0^1 ts\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)dsdt\\
= \quad & \int_0^1 t\left[\frac{2s}{d-c}\frac{\partial f}{\partial t}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)\right|_0^1\\
& -\frac{2}{d-c}\int_0^1\frac{\partial f}{\partial t}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)ds\Big]dt\\
= \quad & \frac{2}{d-c}\int_0^1 t\frac{\partial f}{\partial t}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)dt\\
& -\frac{2}{d-c}\int_0^1\int_0^1 t\frac{\partial f}{\partial t}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)dsdt.
\end{aligned}
$$

By integrating again, we get

$$
\begin{aligned}
& I_1\\
= \quad & \int_0^1\int_0^1 ts\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)dsdt\\
= \quad & \frac{4}{(b-a)(d-c)}\left[f\left(\frac{a+b}{2}, \frac{c+d}{2}\right)-\int_0^1 f\left(t\frac{a+b}{2}+(1-t)a, \frac{c+d}{2}\right)dt\right.\\
& \left.-\int_0^1 f\left(\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)ds+\int_0^1\int_0^1 f\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)dtds\right].
\end{aligned}
$$

Analogously, we can easily obtain the followings

$$
\begin{aligned}
& I_2 \\
= & \int_0^1 \int_0^1 (t-1)(s-1) \frac{\partial^2 f}{\partial t \partial s}\left(tb+(1-t)\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right) dsdt \\
= & \frac{4}{(b-a)(d-c)}\left[f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \int_0^1 f\left(tb+(1-t)\frac{a+b}{2}, \frac{c+d}{2}\right) dt\right. \\
& \left. - \int_0^1 f\left(\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right) ds + \int_0^1 \int_0^1 f\left(tb+(1-t)\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right) dtds\right],
\end{aligned}
$$

$$
\begin{aligned}
& I_3 \\
= & \int_0^1 \int_0^1 t(s-1) \frac{\partial^2 f}{\partial t \partial s}\left(t\frac{a+b}{2}+(1-t)a, sd+(1-s)\frac{c+d}{2}\right) dsdt \\
= & \frac{4}{(b-a)(d-c)}\left[f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \int_0^1 f\left(t\frac{a+b}{2}+(1-t)a, \frac{c+d}{2}\right) dt\right. \\
& \left. - \int_0^1 f\left(\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right) ds + \int_0^1 \int_0^1 f\left(t\frac{a+b}{2}+(1-t)a, sd+(1-s)\frac{c+d}{2}\right) dtds\right]
\end{aligned}
$$

and

$$
\begin{aligned}
& I_4 \\
= & \int_0^1 \int_0^1 (t-1)s \frac{\partial^2 f}{\partial t \partial s}\left(tb+(1-t)\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right) dsdt \\
= & \frac{4}{(b-a)(d-c)}\left[f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \int_0^1 f\left(tb+(1-t)\frac{a+b}{2}, \frac{c+d}{2}\right) dt\right. \\
& \left. - \int_0^1 f\left(\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right) ds + \int_0^1 \int_0^1 f\left(tb+(1-t)\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right) dtds\right].
\end{aligned}
$$

By adding I_1, I_2, I_3, I_4 and changing of the variables, we get the desired result. □

Theorem 2.2 *Let $f : \Delta = [a,b]\times[c,d] \to \mathbb{R}$ be a partial differentiable mapping on $\Delta = [a,b]\times[c,d]$. If $\left|\frac{\partial^2 f}{\partial t \partial s}\right|$ is co-ordinated convex function on the co-ordinates on Δ, then the following inequality holds;*

$$
\begin{aligned}
& |\Lambda| \qquad (10) \\
\leq & \frac{(b-a)(d-c)}{144}\left[\frac{\left|\frac{\partial^2 f}{\partial t \partial s}(a,c)\right| + \left|\frac{\partial^2 f}{\partial t \partial s}(b,c)\right| + \left|\frac{\partial^2 f}{\partial t \partial s}(a,d)\right| + \left|\frac{\partial^2 f}{\partial t \partial s}(b,d)\right|}{4}\right. \\
& +4\left|\frac{\partial^2 f}{\partial t \partial s}\left(\frac{a+b}{2}, \frac{c+d}{2}\right)\right| + \left|\frac{\partial^2 f}{\partial t \partial s}\left(\frac{a+b}{2}, c\right)\right| \\
& \left. + \left|\frac{\partial^2 f}{\partial t \partial s}\left(\frac{a+b}{2}, d\right)\right| + \left|\frac{\partial^2 f}{\partial t \partial s}\left(a, \frac{c+d}{2}\right)\right| + \left|\frac{\partial^2 f}{\partial t \partial s}\left(b, \frac{c+d}{2}\right)\right|\right].
\end{aligned}
$$

Proof From Lemma 1 and by using the absolute value for the both sides of the equality, we can write

$$
\begin{aligned}
&|\Lambda| \\
&= \frac{(b-a)(d-c)}{16}\left[\int_0^1\int_0^1 ts\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)\right|dsdt\right. \\
&+\int_0^1\int_0^1 |(t-1)(s-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right)\right|dsdt \\
&+\int_0^1\int_0^1 |t(s-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, sd+(1-s)\frac{c+d}{2}\right)\right|dsdt \\
&\left.+\int_0^1\int_0^1 |s(t-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)\right|dsdt\right].
\end{aligned}
$$

If we use the co-ordinated convexity of $\left|\frac{\partial^2 f}{\partial t\partial s}\right|$, we have

$$
\begin{aligned}
&|\Lambda| \\
&= \frac{(b-a)(d-c)}{16}\left[\int_0^1\int_0^1 ts\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)\right|\right.\right. \\
&\left.+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}\left(a, s\frac{c+d}{2}+(1-s)c\right)\right|\right\}dsdt \\
&+\int_0^1\int_0^1 |(t-1)(s-1)|\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}\left(b, sd+(1-s)\frac{c+d}{2}\right)\right|\right. \\
&\left.+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right)\right|dsdt\right\} \\
&+\int_0^1\int_0^1 |t(s-1)|\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right)\right|dsdt\right. \\
&\left.+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}\left(a, sd+(1-s)\frac{c+d}{2}\right)\right|dsdt\right\} \\
&+\int_0^1\int_0^1 |s(t-1)|\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}\left(b, s\frac{c+d}{2}+(1-s)c\right)\right|\right. \\
&\left.\left.+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)\right|dsdt\right\}\right].
\end{aligned}
$$

By computing these integrals, we obtain

$$
\begin{aligned}
&|\Lambda| \\
&= \frac{(b-a)(d-c)}{16}\left[\frac{1}{3}\int_0^1 s\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)\right|ds\right. \\
&+\frac{1}{6}\int_0^1 s\left|\frac{\partial^2 f}{\partial t\partial s}\left(a, s\frac{c+d}{2}+(1-s)c\right)\right|ds \\
&+\frac{1}{6}\int_0^1 |s-1|\left|\frac{\partial^2 f}{\partial t\partial s}\left(b, sd+(1-s)\frac{c+d}{2}\right)\right|ds \\
&+\frac{1}{3}\int_0^1 |s-1|\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right)\right|ds \\
&+\frac{1}{3}\int_0^1 |s-1|\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right)\right|ds \\
&+\frac{1}{6}\int_0^1 |s-1|\left|\frac{\partial^2 f}{\partial t\partial s}\left(a, sd+(1-s)\frac{c+d}{2}\right)\right|ds \\
&+\frac{1}{6}\int_0^1 s\left|\frac{\partial^2 f}{\partial t\partial s}\left(b, s\frac{c+d}{2}+(1-s)c\right)\right|ds \\
&\left.+\frac{1}{3}\int_0^1 s\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)\right|ds\right].
\end{aligned}
$$

By using co-ordinated convexity of $\left|\frac{\partial^2 f}{\partial t\partial s}\right|$ again and computing the integrals, we get desired result.

□

Remark 2.3 *Suppose that all the assumptions of Theorem 8 are satisfied. If we choose* $\frac{\partial^2 f}{\partial t\partial s}$ *is bounded, i.e.,*

$$\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty = \sup_{(t,s)\in(0,1)\times(0,1)}\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right| < \infty,$$

we get

$$|\Lambda| \le \frac{(b-a)(d-c)}{16}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty$$

which is the inequality (6).

Theorem 2.4 *Let* $f:\Delta=[a,b]\times[c,d]\to\mathbb{R}$ *be a partial differentiable mapping on* $\Delta=[a,b]\times[c,d]$.

If $\left|\frac{\partial^2 f}{\partial t \partial s}\right|^q$ *is a convex function on the co-ordinates on* Δ*, then the following inequality holds;*

$$
\begin{aligned}
&|\Lambda| \qquad (11)\\
\leq\ & \frac{(b-a)(d-c)}{16(p+1)^{\frac{2}{p}}}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},d\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,\frac{c+d}{2}\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{c+d}{2}\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},c\right)\right|}{2}\right.\\
&\left.\times\left(\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right|^q+\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4}\right)^{\frac{1}{q}}\right.
\end{aligned}
$$

where $q>1$.

Proof From Lemma 1, we have

$$
\begin{aligned}
&|\Lambda|\\
=\ & \frac{(b-a)(d-c)}{16}\left[\int_0^1\int_0^1 ts\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a,s\frac{c+d}{2}+(1-s)c\right)\right|dsdt\right.\\
&+\int_0^1\int_0^1|(t-1)(s-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2},sd+(1-s)\frac{c+d}{2}\right)\right|dsdt\\
&+\int_0^1\int_0^1|t(s-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a,sd+(1-s)\frac{c+d}{2}\right)\right|dsdt\\
&\left.+\int_0^1\int_0^1|s(t-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2},s\frac{c+d}{2}+(1-s)c\right)\right|dsdt\right].
\end{aligned}
$$

By applying the well-known Hölder inequality for double integrals, then one has

$$
\begin{aligned}
&|\Lambda| \qquad (12)\\
=\ & \frac{(b-a)(d-c)}{16}\left[\left(\int_0^1\int_0^1(ts)^p\,dsdt\right)^{\frac{1}{p}}\right.\\
&\times\left(\int_0^1\int_0^1\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a,s\frac{c+d}{2}+(1-s)c\right)\right|^q dsdt\right)^{\frac{1}{q}}\\
&+\left(\int_0^1\int_0^1|(t-1)(s-1)|^p\,dsdt\right)^{\frac{1}{p}}\\
&\times\left(\int_0^1\int_0^1\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2},sd+(1-s)\frac{c+d}{2}\right)\right|^q dsdt\right)^{\frac{1}{q}}\\
&\left.+\left(\int_0^1\int_0^1|t(s-1)|^p\,dsdt\right)^{\frac{1}{p}}\right.
\end{aligned}
$$

$$\times\left(\int_0^1\int_0^1\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)\,a,sd+(1-s)\,\frac{c+d}{2}\right)\right|^q dsdt\right)^{\frac{1}{q}}$$

$$+\left(\int_0^1\int_0^1|s\,(t-1)|^p\,dsdt\right)^{\frac{1}{p}}$$

$$\times\left(\int_0^1\int_0^1\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\,\frac{a+b}{2},s\frac{c+d}{2}+(1-s)\,c\right)\right|^q dsdt\right)^{\frac{1}{q}}\Bigg].$$

Since $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$ is a co-ordinated convex function on Δ, we can write for all $(t,s)\in[0,1]\times[0,1]$

$$\begin{aligned}&\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)\,a,s\frac{c+d}{2}+(1-s)\,c\right)\right|^q \qquad (13)\\ \leq\;& ts\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right|^q\\ &+t\,(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},c\right)\right|^q\\ &+(1-t)\,s\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,\frac{c+d}{2}\right)\right|^q\\ &+(1-t)\,(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\,(a,c)\right|^q.\end{aligned}$$

Similarly, we have

$$\begin{aligned}&\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\,\frac{a+b}{2},sd+(1-s)\,\frac{c+d}{2}\right)\right|^q \qquad (14)\\ \leq\;& ts\left|\frac{\partial^2 f}{\partial t\partial s}\,(b,d)\right|^q\\ &+t\,(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{c+d}{2}\right)\right|^q\\ &+(1-t)\,s\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},d\right)\right|^q\\ &+(1-t)\,(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right|^q,\end{aligned}$$

$$\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)\,a, sd+(1-s)\,\frac{c+d}{2}\right)\right|^q \tag{15}$$

$$\begin{aligned}\leq\ & ts\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, d\right)\right|^q \\ & +t\,(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, \frac{c+d}{2}\right)\right|^q \\ & +(1-t)\,s\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q \\ & +(1-t)\,(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(a, \frac{c+d}{2}\right)\right|^q,\end{aligned}$$

$$\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\,\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)\,c\right)\right|^q \tag{16}$$

$$\begin{aligned}\leq\ & ts\left|\frac{\partial^2 f}{\partial t\partial s}\left(b, \frac{c+d}{2}\right)\right|^q \\ & +t\,(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q \\ & +(1-t)\,s\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, \frac{c+d}{2}\right)\right|^q \\ & +(1-t)\,(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, c\right)\right|^q.\end{aligned}$$

Using inequalities of (13)-(16) in (12), we get

$$\begin{aligned}&|\Lambda| \\ \leq\ & \frac{(b-a)\,(d-c)}{16\,(p+1)^{\frac{2}{p}}}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, d\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(a, \frac{c+d}{2}\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(b, \frac{c+d}{2}\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, c\right)\right|}{2}\right. \\ & \left.\times\left(\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, \frac{c+d}{2}\right)\right|^q+\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4}\right)^{\frac{1}{q}}\right.\end{aligned}$$

This completes the proof. □

Remark 2.5 *Suppose that all the assumptions of Theorem 9 are satisfied. If we choose* $\frac{\partial^2 f}{\partial t\partial s}$ *is bounded, i.e.,*

$$\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty = \sup_{(t,s)\in(0,1)\times(0,1)}\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right| < \infty,$$

we get

$$|\Lambda| \leq \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}} \left\| \frac{\partial^2 f(t,s)}{\partial t \partial s} \right\|_{\infty}$$

which is the inequality (8).

Theorem 2.6 *Let $f : \Delta = [a,b]\times[c,d] \to \mathbb{R}$ be a partial differentiable mapping on $\Delta = [a,b]\times[c,d]$. If $\left|\frac{\partial^2 f}{\partial t \partial s}\right|^q$ is a convex function on the co-ordinates on Δ, then the following inequality holds;*

$$\begin{aligned} &|\Lambda| \\ \leq\ & \frac{(b-a)(d-c)}{16}\left[\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{36}\right. \\ &\left. + \frac{4\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},c\right)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},d\right)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}\left(a,\frac{c+d}{2}\right)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{c+d}{2}\right)\right|^q}{9}\right]^{\frac{1}{q}}. \end{aligned}$$

Proof From Lemma 1, we have

$$\begin{aligned} &|\Lambda| \\ =\ & \frac{(b-a)(d-c)}{16}\left[\int_0^1\int_0^1 ts\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)\right|dsdt\right. \\ &+\int_0^1\int_0^1 |(t-1)(s-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right)\right|dsdt \\ &+\int_0^1\int_0^1 |t(s-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, sd+(1-s)\frac{c+d}{2}\right)\right|dsdt \\ &\left.+\int_0^1\int_0^1 |s(t-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)\right|dsdt\right]. \end{aligned}$$

By applying the well-known Power mean inequality for double integrals, then one has

$$
\begin{aligned}
&|\Lambda| \qquad (17)\\
&= \frac{(b-a)(d-c)}{16}\left[\left(\int_0^1\int_0^1 (ts)\,dsdt\right)^{1-\frac{1}{q}}\right.\\
&\times\left(\int_0^1\int_0^1 (ts)\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)\right|^q dsdt\right)^{\frac{1}{q}}\\
&+\left(\int_0^1\int_0^1 |(t-1)(s-1)|\,dsdt\right)^{1-\frac{1}{q}}\\
&\times\left(\int_0^1\int_0^1 |(t-1)(s-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right)\right|^q dsdt\right)^{\frac{1}{q}}\\
&+\left(\int_0^1\int_0^1 |t(s-1)|\,dsdt\right)^{1-\frac{1}{q}}\\
&\times\left(\int_0^1\int_0^1 |t(s-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, sd+(1-s)\frac{c+d}{2}\right)\right|^q dsdt\right)^{\frac{1}{q}}\\
&+\left(\int_0^1\int_0^1 |s(t-1)|\,dsdt\right)^{1-\frac{1}{q}}\\
&\left.\times\left(\int_0^1\int_0^1 |s(t-1)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)\right|^q dsdt\right)^{\frac{1}{q}}\right].
\end{aligned}
$$

Since $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$ is a co-ordinated convex function on Δ, we can write for all $(t,s)\in[0,1]\times[0,1]$

$$
\begin{aligned}
&\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, s\frac{c+d}{2}+(1-s)c\right)\right|^q \qquad (18)\\
\le\ & ts\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, \frac{c+d}{2}\right)\right|^q\\
&+t(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2}, c\right)\right|^q\\
&+(1-t)s\left|\frac{\partial^2 f}{\partial t\partial s}\left(a, \frac{c+d}{2}\right)\right|^q\\
&+(1-t)(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q.
\end{aligned}
$$

$$\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, sd+(1-s)\frac{c+d}{2}\right)\right|^q \tag{19}$$

$$\begin{aligned}\leq\ & ts\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q \\ & +t(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{c+d}{2}\right)\right|^q \\ & +(1-t)s\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},d\right)\right|^q \\ & +(1-t)(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right|^q,\end{aligned}$$

$$\left|\frac{\partial^2 f}{\partial t\partial s}\left(t\frac{a+b}{2}+(1-t)a, sd+(1-s)\frac{c+d}{2}\right)\right|^q \tag{20}$$

$$\begin{aligned}\leq\ & ts\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},d\right)\right|^q \\ & +t(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right|^q \\ & +(1-t)s\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q \\ & +(1-t)(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,\frac{c+d}{2}\right)\right|^q,\end{aligned}$$

$$\left|\frac{\partial^2 f}{\partial t\partial s}\left(tb+(1-t)\frac{a+b}{2}, s\frac{c+d}{2}+(1-s)c\right)\right|^q \tag{21}$$

$$\begin{aligned}\leq\ & ts\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{c+d}{2}\right)\right|^q \\ & +t(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q \\ & +(1-t)s\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right|^q \\ & +(1-t)(1-s)\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},c\right)\right|^q.\end{aligned}$$

If we use (18)-(21) in (17), we get

$$
\begin{aligned}
&|\Lambda| \\
\leq\ & \frac{(b-a)(d-c)}{16}\left[\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{36}\right. \\
&\left.+\frac{4\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},c\right)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{a+b}{2},d\right)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,\frac{c+d}{2}\right)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{c+d}{2}\right)\right|^q}{9}\right]^{\frac{1}{q}}.
\end{aligned}
$$

This completes the proof. □

References

[1] Bakula M.K., Pečarić J., *On the Jensen's inequality for convex functions on the co-ordinates in a rectangle from the plane*, Taiwanese Journal of Mathematics, 10 (5), 1271-1292, 2006.

[2] Özdemir M.E., Set E., Sarıkaya M.Z., *Some new Hadamard's type inequalities for co-ordinated $m-$convex and $(\alpha,m)-$convex functions*, Hacettepe Journal of Mathematics and Statistics, 40, 219-229, 2011.

[3] Özdemir M.E., Latif M.A., Akdemir A.O., *On some Hadamard-type inequalities for product of two $s-$convex functions on the co-ordinates*, Journal of Inequalities and Applications, 21, 2012.

[4] Set E., Sarıkaya M.Z., Akdemir A.O., *A new general inequality for double integrals*, American Institu of Phsyics (AIP) Conference Proceedings, 1470, 122-125, 2012.

[5] Alomari M., Darus M., *Hadamard-type inequalities for $s-$convex functions*, International Mathematical Forum, 40(3), 1965-1975, 2008.

[6] Özdemir M.E., A.O. Akdemir, *On the hadamard type inequalities involving product of two convex functions on the co-ordinates*, Tamkang Journal of Mathematics, 46(2), 129-142, 2015.

[7] Akdemir A.O., Özdemir M.E., *Some Hadamard-type inequalities for co-ordinated $P-$convex functions and Godunova-Levin functions*, American Institu of Phsyics (AIP) Conference Proceedings, 1309, 7-15, 2010.

[8] Özdemir M.E., Latif M.A., Akdemir A.O., *On some Hadamard-type inequalities for product of two $h-$convex functions on the co-ordinates*, Turkish Journal of Science, 1(1), 48-58, 2016.

[9] Özdemir M.E., Akdemir A.O., Yıldız C., *On co-ordinated quasi-convex functions*, Czechoslovak Mathematical Journal, 62(4), 889-900, 2012.

[10] Alomari M., Darus M., *The Hadamard's inequality for $s-$convex functions of $2-$variables*, International Journal of Mathematical Analysis, 2(13), 629-638, 2008.

[11] Sarıkaya M.Z., Set E., Özdemir M.E., Dragomir S.S., *New some Hadamard's type inequalities for co-ordinated convex functions*, Tamsui Oxford Journal of Information and Mathematical Sciences, 28(2), 137-152, 2012.

[12] Özdemir M.E., Yıldız C., Akdemir A.O., *On some new Hadamard-type inequalities for co-ordinated quasi-convex functions*, Hacettepe Journal of Mathematics and Statistics, 41(5), 697-707, 2012.

[13] Dragomir S.S., *On Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane*, Taiwanese Journal of Mathematics, 5, 775-788, 2001.

[14] Hwang D.Y., Tseng K.L., Yang G.S., *Some Hadamard's inequalities for co-ordinated convex functions in a rectangle from the plane*, Taiwanese Journal of Mathematics, 11, 63-73, 2007.

[15] Özdemir M.E., Kavurmacı H., Akdemir A.O., Avcı M., *Inequalities for convex and $s-$ convex functions on $\Delta = [a,b] \times [c,d]$*, Journal of Inequalities and Applications, 20, 2012.

[16] Bakula M.K., Özdemir M.E., Pečarić J., *Hadamard-type inequalities for $m-$ convex and $(\alpha,m)-$ convex functions*, Journal of Inequalities in Pure and Applied Mathematics, 9(4), 96, 2008.

[17] Bakula M.K, Pečarić J., Ribibić M., *Companion inequalities to Jensen's inequality for $m-$ convex and $(\alpha,m)-$ convex functions*, Journal of Inequalities in Pure and Applied Mathematics, 7(5), 194, 2006.

[18] Dragomir S.S., Toader G., *Some inequalities for $m-$ convex functions*, Studia University Babes Bolyai Mathematica, 38(1), 21-28, 1993.

[19] Miheşan V.G., A Generalization of the Convexity, Seminar of Functional Equations, Approx. and Convex, 1993.

[20] Toader G., *Some generalization of the convexity*, Proceedings of the Colloquium on Approximation and Optimization, 329-338, 1984.

[21] Set E., Sardari M., Özdemir M.E., Rooin J., *On generalizations of the Hadamard inequality for $(\alpha,m)-$ convex functions*, Research Group in Mathematical Inequalities and Applications, 12(4), 4, 2009.

[22] Özdemir M.E., Avcı M., Set E., *On some inequalities of Hermite-Hadamard type via $m-$ convexity*, Applied Mathematics Letters, 23, 1065-1070, 2010.

[23] Toader G., *On a generalization of the convexity*, Mathematica, 30(53), 83-87, 1988.

[24] Dragomir S.S., *On some new inequalities of Hermite-Hadamard type for $m-$ convex functions*, Tamkang Journal of Mathematics, 33(1), 45-56, 2002.

[25] Set E., Özdemir M.E., Dragomir S.S., *On the Hermite-Hadamard inequality and other integral inequalities involving two functions*, Journal of Inequalities and Applications, ID 148102, 2010.

[26] Özdemir M.E., Kavurmacı H., Set E., *Ostrowski's type inequalities for $(\alpha,m)-$ convex functions*, Kyungpook Mathematical Journal, 50(3), 371-378, 2010.

[27] Özdemir M.E., Avcı M., Kavurmacı H., *Hermite-Hadamard type inequalities via $(\alpha,m)-$ convexity*, Computers & Mathematics with Applications, 61(9), 2614-2620, 2011.