Proc. Natl. Acad. Sci., India, Sect. A Phys. Sci. (January–March 2021) 91(1):79–88
https://doi.org/10.1007/s40010-020-00679-9

RESEARCH ARTICLE

Some Inequalities for the First General Zagreb Index of Graphs and Line Graphs

B. Chaluvaraju[1] · H. S. Boregowda[2] · Ismail Naci Cangul[3]

Received: 6 May 2018 / Revised: 11 March 2020 / Accepted: 13 March 2020 / Published online: 2 April 2020
© The National Academy of Sciences, India 2020

Abstract The first general Zagreb index $M_1^{\alpha}(G)$ of a graph G is equal to the sum of the αth powers of the vertex degrees of G. For $\alpha \geq 0$ and $k \geq 1$, we obtain the lower and upper bounds for $M_1^{\alpha}(G)$ and $M_1^{\alpha}(L(G))$ in terms of order, size, minimum/maximum vertex degrees and minimal non-pendant vertex degree using some classical inequalities and majorization technique, where $L(G)$ is the line graph of G. Also, we obtain some bounds and exact values of $M_1^{\alpha}(J(G))$ and $M_1^{\alpha}(L^k(G))$, where $J(G)$ is a jump graph (complement of a line graph) and $L^k(G)$ is an iterated line graph of a graph G.

Keywords First Zagreb index · General first Zagreb index · Line graph · Jump graph · Iterated line graph

Mathematics Subject Classification 05C05 · 05C07 · 05C35

✉ Ismail Naci Cangul
cangul@uludag.edu.tr

B. Chaluvaraju
bchaluvaraju@gmail.com

H. S. Boregowda
bgsamarasa@gmail.com

[1] Department of Mathematics, Bangalore University, Jnana Bharathi Campus, Bengaluru 560 056, India

[2] Department of Studies and Research in Mathematics, University Constituent College Campus, Tumkur University, Tumkur 571 103, India

[3] Department of Mathematics, Faculty of Arts and Science, Bursa Uludag University, Gorukle Campus Bursa, 16059 Bursa, Turkey

1 Introduction

The graphs considered here are finite, undirected, without loops and multiple edges. Let $G = (V, E)$ be a connected graph with $|V(G)| = n$ vertices and $|E(G)| = m$ edges. The degree $d_G(v)$ of a vertex v is the number of vertices adjacent to v. The edge e connecting the vertices u and v is denoted by $e = uv$. The degree of an edge $e = uv$ is denoted by $d_G(e)$ and similarly defined by $d_G(e) = d_G(u) + d_G(v) - 2$. The vertices and edges of a graph are said to be its elements, [1, 2].

A molecular graph is a graph in which the vertices correspond to the atoms and the edges to the bonds of a molecule. A single number that can be computed from the molecular graph and used to characterize some property of the underlying molecule is said to be a topological index or molecular structure descriptor. Numerous such descriptors have been considered in theoretical chemistry and have found some applications, especially in QSPR/QSAR research, [3, 4].

The first Zagreb index was introduced by Gutman and Trinajstić [5]. It is an important molecular descriptor and has been closely correlated with many chemical properties. The first Zagreb index of G is defined as

$$M_1(G) = \sum_{u \in V(G)} d_G(u)^2 \text{ or}$$

$$M_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)].$$

For their history, applications and mathematical properties, see [6–13] and the references cited therein.

Li et al. [14, 15] introduced the generalized version of the first Zagreb index defined as

$$M_1^{\alpha}(G) = \sum_{u \in V(G)} d_G(u)^{\alpha},$$

where α is an arbitrary real number. This graph invariant is also known as "general zeroth-order Randić index," [16, 17].

Following $M_1(G)$, the forgotten topological index (also called F-index) was introduced by Furtula and Gutman [18] and is defined as

$$F(G) = \sum_{v \in V(G)} d_G(u)^3 = \sum_{uv \in E(G)} \left[d_G(u)^2 + d_G(v)^2\right].$$

Shirdel et al. [19] introduced the first hyper-Zagreb index of G, which is defined as

$$HM_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^2.$$

Clearly, we note that $M_1^0(G) = n$, $M_1^1(G) = 2m$, $M_1^2(G) = M_1(G)$ and $M_1^3(G) = F(G)$ and $M_1^4(G) = HM_1(G)$.

2 Bounds on $M_1^{\alpha}(G)$

First, we prove a lower bound for $M_1^{\alpha}(G)$ in terms of order n and size m. For this purpose, we recall the following facts.

If a real-valued function $f(x)$ defined on an interval has a second derivative $f''(x)$, then a necessary and sufficient condition for it to be strictly convex on that interval is that $f''(x) > 0$. For a positive integer k, if $f(x)$ is strictly convex, then by Jensen's inequality, we have $f\left(\sum_{i=1}^{k} \frac{x_i}{k}\right) \leq \frac{1}{k}\sum_{i=1}^{k} f(x_i)$ with equality if and only if $x_1 = x_2 = \cdots = x_k$, and if $-f(x)$ is strictly convex, then the inequality is reversed.

Theorem 1 *Let G be a simple graph with $n \geq 3$ vertices and m edges. Then, for all $\alpha \geq 1$,*

$$M_1^{\alpha}(G) \geq \frac{2^{\alpha} m^{\alpha}}{n^{\alpha-1}}.$$

Further, equality holds if and only if G is regular.

Proof Let G be a simple graph with $n \geq 3$ vertices and m edges. We have $M_1^{\alpha}(G) = \sum_{v \in V(G)} d_G(v)^{\alpha}$, for all $\alpha \geq 1$. Since $f(x) = x^{\alpha}$ is strictly convex for $x > 0$. By Jensen's inequality, we have

$$\sum_{i=1}^{k} f(x_i) \geq k f\left(\sum_{i=1}^{k} \frac{x_i}{k}\right).$$

Therefore,

$$\begin{aligned} M_1^{\alpha}(G) &\geq n\left(\sum_{u \in V(G)} \frac{d_G(u)}{n}\right)^{\alpha} \\ &\geq n\left(\frac{2m}{n}\right)^{\alpha} \\ &\geq \frac{2^{\alpha} m^{\alpha}}{n^{\alpha-1}}. \end{aligned}$$

□

Corollary 1 *For any (n, m)-simple graph G with $n \geq 3$ vertices and $\alpha = 2$,*

$$M_1^2(G) = M_1(G) \geq \frac{4m^2}{n}.$$

Theorem 2 *Let G be a connected graph with $n \geq 3$ vertices and m edges. Then, for all $\alpha \geq 1$, we have*

$$M_1^{\alpha}(G) \geq n\left(\prod_{u \in V(G)} (d_G(u))^{\alpha}\right)^{\frac{1}{n}}.$$

Proof Let $f(x) = \log x$. Clearly, $\log(x)$ is concave in $(0, \infty)$. By Jensen's inequality, we have

$$\begin{aligned} \log\left(\sum_{u \in V(G)} \frac{(d_G(u))^{\alpha}}{n}\right) &\geq \frac{1}{n}\sum_{u \in V(G)} \log(d_G(u))^{\alpha} \\ &\geq \frac{1}{n}\log\left(\prod_{u \in V(G)} d_G(u)^{\alpha}\right) \\ &\geq \log\left(\prod_{u \in V(G)} d_G(u)^{\alpha}\right)^{\frac{1}{n}}. \end{aligned}$$

Therefore,

$$\frac{1}{n}\sum_{u \in V(G)} d_G(u)^{\alpha} \geq \left(\prod_{u \in V(G)} d_G(u)^{\alpha}\right)^{\frac{1}{n}}$$

$$\sum_{u \in V(G)} (d_G(u))^{\alpha} \geq n\left(\prod_{u \in V(G)} d_G(u)^{\alpha}\right)^{\frac{1}{n}}.$$

Thus, the upper bound follows. □

In order to prove our next result giving lower and upper bounds for $M_1^{\alpha}(G)$ in terms of order n, minimum degree $\delta(G)$ and maximum degree $\Delta(G)$ of a graph G, we apply the well-known Diaz-Metcalf inequality.

Theorem 3 [20] *Let $(a_1, a_2, \ldots, a_n)$ and $(b_1, b_2, \ldots, b_n)$ be two n-tuples consisting of positive real numbers satisfying the condition*

$$ra_i \leq b_i \leq Ra_i$$

for $1 \le i \le n$. *Then,*

$$\sum_{i=1}^{n} b_i^2 + rR \sum_{i=1}^{n} a_i^2 \le (r+R) \sum_{i=1}^{n} a_i b_i.$$

Equality holds if and only if $b_i = Ra_i$ or $b_i = ra_i$ *for* $1 \le i \le n$.

Theorem 4 *Let G be a simple graph with* $n \ge 3$ *vertices and m edges. Then, for all* $\alpha \ge 1$,

$$n\,\delta(G)^{\alpha} \le M_1^{\alpha}(G) \le n\,\Delta(G)^{\alpha}.$$

Further, both left- and right-hand side equality holds if and only if G is regular.

Proof Let $b_i = d_G(u)^{\frac{\alpha}{2}}$; $a_i = 1$ and $r = \delta(G)^{\frac{\alpha}{2}}$; $R = \Delta(G)^{\frac{\alpha}{2}}$. Clearly,

$$\delta(G)^{\frac{\alpha}{2}} \le d_G(u)^{\frac{\alpha}{2}} \le \Delta(G)^{\frac{\alpha}{2}}.$$

By the Diaz–Metcalf inequality, we have

$$\sum_{u \in V(G)} d_G(u)^{a} + (\delta(G)\Delta(G))^{\frac{\alpha}{2}} \sum_{u \in V(G)} 1 \le (\delta(G)^{\frac{\alpha}{2}} + \Delta(G)^{\frac{\alpha}{2}}) \sum_{u \in V(G)} d_G(u)^{\frac{\alpha}{2}},$$

and hence,

$$M_1^{\alpha}(G) + n\,\delta(G)^{\frac{\alpha}{2}} \le (\delta(G)^{\frac{\alpha}{2}} + \Delta(G)^{\frac{\alpha}{2}}) n\,\Delta(G)^{\frac{\alpha}{2}}.$$

Therefore, we have $d_G(u)^{\frac{\alpha}{2}} \le \Delta(G)^{\frac{\alpha}{2}}$ for all $u \in V(G)$. Hence, we can write

$$M_1^{\alpha}(G) \le (\delta(G)^{\frac{\alpha}{2}} + \Delta(G)^{\frac{\alpha}{2}}) n\,\Delta(G)^{\frac{\alpha}{2}} - n\,\delta(G)^{\frac{\alpha}{2}}$$

and

$$M_1^{\alpha}(G) \le n\,\delta(G)^{\frac{\alpha}{2}} + n\,\Delta(G)^{\frac{\alpha}{2}} - n\,\delta(G)^{\frac{\alpha}{2}} = n\,\Delta(G)^{\alpha}.$$

Thus, the upper bound follows.

Let $b_i = d_G(u)^{\alpha}$; $a_i = 1$ and $r = \delta(G)^{\alpha}$; $R = \Delta(G)^{\alpha}$. By the Diaz–Metcalf inequality, we have

$$\sum_{u \in V(G)} d_G(u)^{2\alpha} + (\delta(G)\Delta(G))^{\alpha} \sum_{u \in V(G)} 1 \le (\delta(G)^{\alpha} + \Delta(G)^{\alpha}) \sum_{u \in V(G)} d_G(u)^{\alpha}$$

and

$$\begin{aligned} M_1^{\alpha}(G) &\ge \frac{1}{(\delta(G)^{\alpha} + \Delta(G)^{\alpha})} \left[\sum_{u \in V(G)} d_G(u)^{2a} + n\left(\delta(G)\Delta(G)\right)^{\alpha} \right] \\ &\ge \frac{1}{(\delta(G)^{\alpha} + \Delta(G)^{\alpha})} \left[n\,\delta(G)^{2\alpha} + n\,\delta(G)^{\alpha}\Delta(G)^{\alpha} \right] \\ &\ge \frac{1}{(\delta(G)^{\alpha} + \Delta(G)^{\alpha})}\, n\,\delta(G)^{\alpha}[\delta(G)^{\alpha} + \Delta(G)^{\alpha}] = n\,\delta(G)^{\alpha}. \end{aligned}$$

Thus, the lower bound follows.

Equality is attained if and only if G is regular. □

The following results are immediate from Theorem 4.

Corollary 2 *For any graph G with* $n \ge 3$ *vertices and* $\alpha = 2$,

$$M_1^2(G) = M_1(G) \le n\,\Delta(G)^2.$$

Corollary 3 *For any complete graph* K_n *with* $n \ge 3$ *vertices and* $\alpha = 2$,

$$M_1^2(K_n) = n(n-1)^2.$$

Our next result gives lower and upper bounds for $M_1^{\alpha}(G)$ in terms of size m, minimum degree $\delta(G)$ and maximum degree $\Delta(G)$ of a graph G.

Theorem 5 *Let G be a simple graph with* $n \ge 3$ *vertices and m edges. Then, for all* $\alpha \ge 0$,

$$2m\,\delta(G)^{\alpha-1} \le M_1^{\alpha}(G) \le 2m\,\Delta(G)^{\alpha-1}.$$

Further, both left- and right-hand side equality holds if and only if G is regular.

Proof Let f be a positive function defined on nonnegative integers. Then,

$$\begin{aligned} X(G) &= \sum_{u \in V(G)} f(d_G(u)) = \sum_{u \in V(G)} \sum_{uv \in E(G)} \frac{f(d_G(u))}{d_G(u)} \\ &= \sum_{uv \in E(G)} \left(\frac{f(d_G(u))}{d_G(u)} + \frac{f(d_G(v))}{d_G(v)} \right), \end{aligned}$$

where $X(G)$ is any degree-based topological index of a graph G.

Defining $f(d_G(u)) = (d_G(u))^{\alpha}$, we have

$$M_1^{\alpha}(G) = \sum_{uv \in E(G)} \left[(d_G(u))^{\alpha-1} + (d_G(v))^{\alpha-1} \right].$$

Since for all $u \in V(G)$, $\delta(G) \le d_G(u) \le \Delta(G)$. Hence, we have

$$\sum_{uv \in E(G)} 2\,\delta(G)^{\alpha-1} \le M_1^{\alpha}(G)$$

$$2\,m\,\delta(G)^{\alpha-1} \le M_1^{\alpha}(G)$$

Equality is attained on both sides if and only if G is regular. □

Now, to obtain another result for the lower and upper bounds for $M_1^{\alpha}(G)$ in terms of order n, minimum degree $\delta(G)$ and maximum degree $\Delta(G)$ of a graph G, we apply the Polya–Szego inequality as follows.

Theorem 6 [21] *Let* $a = (a_1, a_2, \ldots, a_n)$ *and* $b = (b_1, b_2, \ldots, b_n)$ *be two n-tuples of positive numbers. If*

$0<\gamma\le a_i\le A<\infty$ *and* $0<\beta\le b_i\le B<\infty$ *for each* $i\in\{1,2,\ldots,n\}$, *then*

$$\sum_{i=1}^{n}a_i^2\cdot\sum_{i=1}^{n}b_i^2\le\frac{(\gamma\beta+AB)^2}{4\gamma\beta AB}\left(\sum_{i=1}^{n}a_ib_i\right)^2.$$

The equality holds iff $p=n\frac{A}{\gamma}\Big/\left(\frac{A}{\gamma}+\frac{B}{\beta}\right)$ *and* $q=n.\frac{B}{\beta}\Big/\left(\frac{A}{\gamma}+\frac{B}{\beta}\right)$ *are integers and* p *of the numbers* $a_1,a_2,\ldots,a_n$ *are equal to* γ *and* q *of these numbers are equal to* A, *and similarly if* q *of the corresponding numbers* b_i *are equal to* B *and and* q *of them are equal to* β.

Theorem 7 *For any* (n, m)*-connected graph* G *with* $n\ge3$ *vertices and* $\alpha\ge0$, *we have*

$$\frac{2n\,\delta(G)^{2\alpha}}{(\delta(G)^{\alpha}+\Delta(G)^{\alpha})}\left(\frac{\Delta(G)}{\delta(G)}\right)^{\frac{\alpha}{2}}\le M_1^{\alpha}(G)\le\frac{n}{4}\left(\frac{\Delta(G)}{\delta(G)}\right)^{\frac{\alpha}{2}}\left(\delta(G)^{\frac{\alpha}{2}}+\Delta(G)^{\frac{\alpha}{2}}\right)^2.$$

Further, both left- and right-hand side equalities hold if and only if G *is regular.*

Proof Let $a_i=d_G(u)^{\frac{\alpha}{2}}$ and $b_i=1$ in the Polya–Szego inequality. Clearly, $\gamma=\delta(G)^{\frac{\alpha}{2}}$, $A=\Delta(G)^{\frac{\alpha}{2}}$, $\beta=1$ and $B=1$. We have

$$\sum_{u\in V(G)}d_G(u)^a\,n\le\frac{\left(\delta(G)^{\frac{\alpha}{2}}+\Delta(G)^{\frac{\alpha}{2}}\right)}{4\,\delta(G)^{\frac{\alpha}{2}}\Delta(G)^{\frac{\alpha}{2}}}\left[\sum_{u\in V(G)}d_G(u)^{\frac{\alpha}{2}}\right]^2\le\frac{\left(\delta(G)^{\frac{\alpha}{2}}+\Delta(G)^{\frac{\alpha}{2}}\right)}{4\,\delta(G)^{\frac{\alpha}{2}}\Delta(G)^{\frac{\alpha}{2}}}\left[n\,\Delta(G)^{\frac{\alpha}{2}}\right]^2.$$

Since $d_G(u)^{\frac{\alpha}{2}}\le\Delta(G)^{\frac{\alpha}{2}}$ for all $u\in V(G)$, we have

$$M_1^{\alpha}(G)\le\frac{\left(\delta(G)^{\frac{\alpha}{2}}+\Delta(G)^{\frac{\alpha}{2}}\right)}{4\,n\,\delta(G)^{\frac{\alpha}{2}}\Delta(G)^{\frac{\alpha}{2}}}\left[n^2\,\Delta(G)^{\alpha}\right]\le\frac{n}{4}\left(\frac{\Delta(G)}{\delta(G)}\right)^{\frac{\alpha}{2}}\left(\delta(G)^{\frac{\alpha}{2}}+\Delta(G)^{\frac{\alpha}{2}}\right)^2.$$

Thus, the upper bound follows.

Let $a_i=d_G(u)^{\alpha}$ and $b_i=1$ in the Polya–Szego inequality. Clearly, $\gamma=\delta(G)^{\alpha}$, $A=\Delta(G)^{\alpha}$, $\beta=1$ and $B=1$. We have

$$M_1^{\alpha}(G)\ge\frac{4\,n\,(\delta(G)\Delta(G))^{\alpha}}{(\delta(G)^{\alpha}+\Delta(G)^{\alpha})^2}\sum_{u\in V(G)}d_G(u)^{2\alpha}\ge$$

Thus, the lower bound follows.

Equality is attained on both sides if and only if G is regular. □

In order to prove our next result giving another upper bound for $M_1^{\alpha}(G)$, this time in terms of the size m of the graph G. We make use of the following well-known majorization technique, [22].

Let $a=(a_1,a_2,\ldots,a_n)$ and $b=(b_1,b_2,\ldots,b_n)$ be two non-increasing n-tuples of real numbers. Then, we say that a is majorized by b denoted by $a\prec b$ if

1. $a_1+a_2+\cdots+a_k\le b_1+b_2+\cdots+b_k$ for $1\le k\le n-1$
2. $a_1+a_2+\cdots+a_n=b_1+b_2+\cdots+b_n$.

Lemma 1 [23] *Suppose* $a=(a_1,a_2,\ldots,a_n)$ *and* $b=(b_1,b_2,\ldots,b_n)$ *are two non-increasing* n*-tuples of real numbers. If* $a\prec b$, *then for any convex function* f,

$$\sum_{j=1}^{n}f(a_j)\le\sum_{j=1}^{n}f(b_j).$$

Theorem 8 *For any* (n, m)*-connected graph* G *with* $n\ge3$ *vertices and for all* $\alpha>1$ *or* $\alpha<0$, *we have*

$$M_1^{\alpha}(G)\le2^{\alpha}\,m^{\alpha}.$$

Proof We prove this result by means of the majorization technique. Clearly, $\left(d_G(u_1),d_G(u_2),\ldots,d_G(u_n)\right)\prec(2m,0,\ldots,0)$ and $f(x)=x^{\alpha}$ is convex for all $\alpha>1$ or $\alpha<0$ and $x>0$. By the above lemma, we have

$$\left(d_G(u_1)^{\alpha}+d_G(u_2)^{\alpha}+\cdots+d_G(u_n)^{\alpha}\right)\le((2m)^{\alpha}+0+\cdots+0),$$

and hence,

$$\sum_{u\in V(G)}d_G(u)^{\alpha}\le(2m)^{\alpha}.$$

Thus, the upper bound follows. □

Corollary 4 *For any* (n, m)*-connected graph* G *with* $n\ge3$ *and* $\alpha=2$,

$$M_1^2(G)\le4\,m^2.$$

To prove our next result, we make use of the following definitions and observations:

A vertex $v\in V(T)$ is called a branching vertex of a tree T, if $\deg_T(v)\ge3$. If $\deg_T(v)=1$, the vertex v is named a leaf (or a pendent vertex) of T. Further, the path with n vertices, of which exactly two are leaves, is denoted by P_n, and the star graph with exactly $n-1$ leaves and one branching vertex is denoted by S_n or $K_{1,n-1}$; $n\ge2$ vertices.

It is easy to see that every tree on n vertices has at least two leaves and at most $\frac{n-2}{2}$ branching vertices.

Theorem 9 *For any nontrivial tree* T *of order* n *with* $\alpha\ge1$,

$$M_1^{\alpha}(P_n)\le M_1^{\alpha}(T)\le M_1^{\alpha}(S_n).$$

Proof We prove the result by using the majorization

technique. Clearly, for the degree sequence of any tree T, we have

$$\big(d_T(u_1), d_T(u_2), \ldots, d_T(u_n)\big) \prec (n-1, \underbrace{1, 1, \ldots, 1}_{(n-1)-\text{times}}).$$

Let $f(x) = x^\alpha$. Clearly, $f(x)$ is convex in $(0, \infty)$. Then,

$$\sum_{u \in V(T)} d_T(u)^\alpha \le (n-1)^\alpha + \underbrace{1 + 1 + \ldots + 1}_{(n-1)-\text{times}} \le (n-1) + (n-1)^\alpha.$$

But, for a tree, we have $m = n-1$, and therefore, we have $M_1^\alpha(T) \le m + m^\alpha$. Thus, the upper bound follows.

Now, the degree sequence of any tree T satisfies

$$\big(1, 2, 2, \ldots 2, 1\big) \prec \big(d_T(u_1), d_T(u_2), \ldots, d_T(u_n)\big).$$

Therefore,

$$1^\alpha + 2^\alpha + \cdots + 2^\alpha + 1^\alpha \le \sum_{u \in V(T)} d_T(u)^\alpha,$$

where $f(x) = x^\alpha$ is convex for all $\alpha \ge 1$. Hence,

$$M_1^\alpha(P_n) = 2 + (n-2)2^\alpha \le M_1^\alpha(T).$$

Thus, the lower bound follows. □

Now, we obtain some lower and upper bounds for $M_1^\alpha(G)$ in terms of the number of pendent vertices and minimal non-pendent vertices of G.

Theorem 10 *For any (n, m)-connected graph G with η-pendent vertices, minimal non-pendent vertex degree $\delta_1(G)$ and $\alpha \ge 0$,*

$$\eta + (m-\eta)\,\delta_1(G)^\alpha \le M_1^\alpha(G) \le \eta + (m-\eta)\,\Delta(G)^\alpha.$$

Further, equality in both lower and upper bounds are attained if and only if G is regular.

Proof For any $\alpha \ge 0$, we have

$$\begin{aligned} M_1^\alpha(G) &= \sum_{u \in V(G)} [d_G(u)]^\alpha \\ &= \sum_{u \in V(G); d_G(u)=1} 1^\alpha + \sum_{u \in V(G); d_G(u) \ne 1} [d_G(u)]^\alpha \\ &\le \eta + (m-\eta)\,\Delta(G)^\alpha. \end{aligned}$$

Similarly,

$$M_1^\alpha(G) \ge \eta + (m-\eta)\,\delta_1(G)^\alpha.$$

Hence, the desired result follows.

Further, the equality in both lower and upper bounds is attained if and only if G is regular. □

Nordhaus and Gaddum [24] gave tight bounds on the sum and product of the chromatic numbers of a graph and its complement. Since then, such type of results have been derived for several other graph invariants. Here, we derive such a relation for $M_1^\alpha(G)$.

Theorem 11 *For any (n, m)-connected graph G with $\alpha \ge 2$ and $n \ge 4$ vertices having a connected $\overline{G}$, we have*

1. $2^{1-\alpha}\, n\,(n-1)^\alpha \le M_1^\alpha(G) + M_1^\alpha(\overline{G}) \le n\,(n-1)^\alpha$,
2. $n^2 \le M_1^\alpha(G) \times M_1^\alpha(\overline{G}) \le n^2 (n-1)^{2\alpha}$.

Proof

1. Since $m + \overline{m} = \frac{n(n-1)}{2}$, we have $d_G(u) + d_{\overline{G}}(u) = n-1$ and $d_G(v) + d_{\overline{G}}(v) = n-1$. Hence, we have

$$\begin{aligned} M_1^\alpha(G) + M_1^\alpha(\overline{G}) &= \sum_{uv \in E(G)} d_G(u)^{\alpha-1} + d_G(v)^{\alpha-1} \\ &\quad + \sum_{uv \notin E(\overline{G})} d_{\overline{G}}(u)^{\alpha-1} + d_{\overline{G}}(v)^{\alpha-1} \\ &\le m\left[2(n-1)^{\alpha-1}\right] + \overline{m}\left[2(n-1)^{\alpha-1}\right] \\ &\le 2(m+\overline{m})\,(n-1)^{\alpha-1}. \end{aligned}$$

Thus, the upper bound follows. By Theorem 1, we have $M_1^\alpha(G) \ge \frac{2^\alpha m^\alpha}{n^{\alpha-1}}$ and $M_1^\alpha(\overline{G}) \ge \frac{2^\alpha(\overline{m})^\alpha}{n^{\alpha-1}}$. Therefore,

$$M_1^\alpha(G) + M_1^\alpha(\overline{G}) \ge \frac{2^\alpha}{n^{\alpha-1}}\,[m^\alpha + \overline{m}^\alpha].$$

Since m^α is convex for all $\alpha \ge 2$, then by Jensen's inequality, we have

$$\begin{aligned} M_1^\alpha(G) + M_1^\alpha(\overline{G}) &\ge \frac{2^\alpha}{n^{\alpha-1}}\; 2\left(\frac{m+\overline{m}}{2}\right)^\alpha \\ &\ge \frac{2^{\alpha+1}}{n^{\alpha-1}}\;\frac{n^\alpha\,(n-1)^\alpha}{2^{2\alpha}}. \end{aligned}$$

Thus, the lower bound follows.

2. Since $1 \le \delta(G) \le \Delta(G) \le n-1$ and $1 \le \delta(\overline{G}) \le \Delta(\overline{G}) \le n-1$ for both G and $\overline{G}$ are connected with $n \ge 4$ vertices, by Theorem 4, we have the desired results.

□

3 Line Graphs

The line graph $L(G)$ is the graph with vertex set $V(L(G)) = E(G)$, and whose vertices correspond to the edges of G with two vertices being adjacent if and only if the corresponding edges in G have a vertex in common. As $L(G)$ is defined on the edge set of a graph G, isolated vertices of G, if there is any, play no role in $L(G)$, [8, 25, 26].

Ranjini et al. [27] obtained the first and second Zagreb indices of the line graphs of the subdivision graphs. Here, we obtain some lower and upper bounds for $M_1^\alpha(L(G))$ in

terms of the size m, minimum degree $\delta(G)$ and maximum degree $\Delta(G)$ of a graph G.

Theorem 12 *Let G be a simple connected graph with $n \geq 3$ vertices and m edges. Then, for all $\alpha \geq 0$, we have*

$$2^{\alpha} m \left(\delta(G) - 1\right)^{\alpha} \leq M_1^{\alpha}(L(G)) \leq 2^{\alpha} m \left(\Delta(G) - 1\right)^{\alpha}.$$

Further, both left- and right-hand side equalities holds if and only if G is regular.

Proof Let G be a simple connected (n, m)-graph with $\alpha \geq 0$. Then,

$$M_1^{\alpha}(L(G)) = \sum_{uv \in E(G)} [d_G(u) + d_G(v) - 2]^{\alpha}.$$

Since for all $u, v \in V(G)$, $\delta(G) \leq \{d_G(u), d_G(v)\} \leq \Delta(G)$, we have

$$\begin{aligned} \sum_{uv \in E(G)} (\delta(G) + \delta(G) - 2)^{\alpha} &\leq M_1^{\alpha}(L(G)) \\ &\leq \sum_{uv \in E(G)} (\Delta(G) + \Delta(G) - 2)^{\alpha} \end{aligned},$$

and therefore,

$$2^{\alpha} m \left(\delta(G) - 1\right)^{\alpha} \leq M_1^{\alpha}(L(G))$$

Further, both left- and right-hand side equalities hold if and only if G is regular. □

We now give our next result giving a lower bound for $M_1^{\alpha}(L(G))$ in terms of the size m and the first Zagreb index $M_1(G)$.

Theorem 13 *Let G be an (n, m)-connected graph with $n \geq 3$ vertices. Then, for all $\alpha \geq 1$, we have*

$$M_1^{\alpha}(L(G)) \geq \frac{\left(M_1(G) - 2m\right)^{\alpha}}{m^{\alpha - 1}}.$$

Further, equality holds if and only if G is regular.

Proof Let G be a (n, m)-connected graph with $n \geq 3$ vertices and m edges. Clearly, $|V(L(G))| = m$ and $|E(L(G))| = \frac{1}{2} M_1(G) - m$. Hence, we have $M_1^{\alpha}(L(G)) = \sum_{v \in V(L(G))} d_{L(G)}(v)^{\alpha}$. For all $\alpha \geq 1$ and $x > 0$, $f(x) = x^{\alpha}$ is strictly convex. Hence, by Jensen's inequality, we have

$$\begin{aligned} M_1^{\alpha}(L(G)) &\geq m \left(\sum_{u \in V(L(G))} \frac{d_{L(G)}(u)}{n} \right)^{\alpha} \\ &\geq m \left(\frac{1}{m} 2 \left(\frac{1}{2} M_1(G) - m \right) \right)^{\alpha} \\ &\geq \frac{\left(M_1(G) - 2m\right)^{\alpha}}{m^{\alpha - 1}}. \end{aligned}$$

Further, equality holds if and only if G is regular. □

Corollary 5 *Let G be an r-regular graph with $n \geq 3$ vertices. Then, for all $\alpha \geq 1$, we have*

$$M_1^{\alpha}(L(G)) = 2^{\alpha - 1} n r (r - 1)^{\alpha}.$$

Let G be an (n, m)-connected graph. Then, $G \cong L(G)$ if and only if $G \cong C_n$; $n \geq 3$. Hence, we have

Corollary 6 *For any cycle C_n with $n \geq 3$ vertices,*

$$M_1^{\alpha}(C_n) = M_1^{\alpha}(L(C_n)) = n 2^{\alpha}.$$

The following is an inequality for $M_1^{\alpha}(L(G))$ in terms of $M_1^{\alpha - i}(L(G))$'s.

Theorem 14 *For any (n, m)-connected graph G with $n \geq 3$ vertices and $\alpha \geq 0$,*

$$M_1^{\alpha}(L(G)) \geq \sum_{i=0}^{\alpha} \binom{\alpha}{i} (-1)^i 2^i m^{-a+i+1} (M_1^{\alpha - i}(G)).$$

Further, equality holds if and only if G is regular.

Proof Let G be a (n, m)-connected graph with $n \geq 3$ and $\alpha \geq 0$. Then,

$$\begin{aligned} M_1^{\alpha}(L(G)) &= \sum_{uv \in E(G)} [d_G(u) + d_G(v) - 2]^{\alpha} \\ &= \sum_{i=0}^{\alpha} \binom{\alpha}{i} (-1)^i \left[d_G(u) + d_G(v)\right]^{\alpha - i} 2^i \\ &= \sum_{i=0}^{\alpha} \binom{\alpha}{i} (-1)^i 2^i \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^{\alpha - i} \\ &= (-1)^{\alpha} 2^{\alpha} m \\ &\quad + \sum_{i=0}^{\alpha - 1} \binom{\alpha}{i} (-1)^i 2^i \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^{\alpha - i}. \end{aligned}$$

By Jensen's inequality as x^{α} is a convex function for $\alpha \geq 1$ and $x > 0$, we have

$$\begin{aligned} M_1^{\alpha}(L(G)) &\geq (-1)^{\alpha} 2^{\alpha} m \\ &\quad + \sum_{i=0}^{\alpha - 1} \binom{\alpha}{i} (-1)^i 2^i m \\ &\quad \sum_{uv \in E(G)} \left[\frac{d_G(u) + d_G(v)}{m} \right]^{\alpha - i} \\ &\geq (-1)^{\alpha} 2^{\alpha} m \\ &\quad + \sum_{i=0}^{\alpha - 1} \binom{\alpha}{i} (-1)^i 2^i m^{i - \alpha + 1} [M_1^{\alpha - 1}(G)] \\ &\geq \sum_{i=0}^{\alpha} \binom{\alpha}{i} (-1)^i 2^i m^{-a+i+1} (M_1^{\alpha - i}(G)). \end{aligned}$$

Further, equality holds if and only if G is regular. □

Corollary 7 *For any connected graph G with $n \geq 3$ vertices and $\alpha = 2$, we have*

$$M_1^2(L(G)) = M_1(L(G)) \geq \frac{(M_1(G))^2}{m} - M_1(G) + 4m.$$

Theorem 15 *Let G be a r-regular connected graph with $\alpha \geq 0$. Then,*

$$M_1^{\alpha}(L(G)) = \sum_{i=0}^{\alpha} \sum_{j=0}^{\alpha-i} \binom{\alpha}{i} \binom{\alpha - i}{j} (-1)^i (2)^{i-1} \; M_1^{\alpha-i+1}(G).$$

Proof Let G be an r-regular connected graph with $\alpha \geq 0$. Then,

$$\begin{aligned} M_1^{\alpha}(L(G)) &= \sum_{uv \in E(G)} [d_G(u) + d_G(v) - 2]^{\alpha} \\ &= \sum_{i=0}^{\alpha} \binom{\alpha}{i} (-1)^i 2^i \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^{\alpha - i} \\ &= \sum_{i=0}^{\alpha} \binom{\alpha}{i} (-1)^i 2^i \sum_{j=0}^{\alpha-i} \binom{\alpha - i}{j} \\ &\quad \sum_{uv \in E(G)} (d_G(u))^{\alpha-i-j} \; (d_G(v))^j. \end{aligned}$$

Since G is an r-regular connected graph as $d_G(u) = d_G(v) = r$, we have

$$\begin{aligned} M_1^{\alpha}(L(G)) &= \sum_{i=0}^{\alpha} \binom{\alpha}{i} (-1)^i 2^i \sum_{j=0}^{\alpha-i} \binom{\alpha - i}{j} \sum_{uv \in E(G)} (d_G(u))^{\alpha - i} \\ &= \sum_{i=0}^{\alpha} \sum_{j=0}^{\alpha-i} \binom{\alpha}{i} (-1)^i 2^i \binom{\alpha - i}{j} \; \frac{1}{2} \sum_{u \in V(G)} (d_G(u))^{\alpha-i+1} \\ &= \sum_{i=0}^{\alpha} \sum_{j=0}^{\alpha-i} \binom{\alpha}{i} \binom{\alpha - i}{j} (-1)^i 2^{i-1} \; M_1^{\alpha-i+1}(G). \end{aligned}$$

□

Corollary 8 *Let G be an r-regular connected graph with $\alpha = 2$. Then,*

$$\begin{aligned} &M_1^2(L(G)) = M_1(L(G)) \\ &= \sum_{i=0}^{2} \sum_{j=0}^{2-i} \binom{2}{i} \binom{2 - i}{j} (-1)^i 2^{i-1} \; M_1^{3-i}(G) \\ &= 2nr(r-1)^2. \end{aligned}$$

In order to prove our next result on $M_1^{\alpha}(L(G))$, we make use of the following trinomial theorem. For more details, we refer to [28].

Theorem 16 *Let x, y, z be three real numbers and n be a whole number. Then,*

$$(x + y + z)^n = \sum_{i,j,k \geq 0; i+j+k=n} \frac{n!}{i!\, j!\, k!} \, x^i y^j z^k.$$

Theorem 17 *Let G be an r-regular connected graph and let α be a whole number. Then,*

$$M_1^{\alpha}(L(G)) = \frac{1}{2} \sum_{i,j,k \geq 0; i+j+k=\alpha} \frac{\alpha!}{i!\, j!\, k!} \, (-2)^k \; M_1^{i+j+1}(G).$$

Proof Let G be an r-regular connected graph and let α be a whole number. Then,

$$\begin{aligned} M_1^{\alpha}(L(G)) &= \sum_{uv \in E(G)} [d_G(u) + d_G(v) - 2]^{\alpha} \\ &= \sum_{uv \in E(G)} \\ &\quad \sum_{i,j,k \geq 0; i+j+k=\alpha} \frac{\alpha!}{i!\, j!\, k!} \, (d_G(u))^i \, (d_G(v))^j \, (-2)^k \\ &= \sum_{i,j,k \geq 0; i+j+k=\alpha} \frac{\alpha!}{i!\, j!\, k!} \, (-2)^k \\ &\quad \sum_{uv \in E(G)} (d_G(u))^i \, (d_G(v))^j. \end{aligned}$$

Since G is r-regular, $d_G(u) = d_G(v) = r$, and hence, we have

$$\begin{aligned} M_1^{\alpha}(L(G)) &= \sum_{i,j,k \geq 0; i+j+k=\alpha} \frac{\alpha!}{i!\, j!\, k!} \, (-2)^k \, \frac{1}{2} \sum_{u \in V(G)} [d_G(u)]^{i+j+1} \\ &= \frac{1}{2} \sum_{i,j,k \geq 0; i+j+k=\alpha} \frac{\alpha!}{i!\, j!\, k!} \, (-2)^k \; M_1^{i+j+1}(G). \end{aligned}$$

□

The following results are immediate from the above theorem.

Corollary 9 *For any r-regular connected graph G with $\alpha = 2$, we have*

$$\begin{aligned} M_1^2(L(G)) = M_1(L(G)) &= \frac{1}{2} \sum_{i,j,k \geq 0; i+j+k=2} \frac{2!}{i!\, j!\, k!} \, (-2)^k \; M_1^{i+j+1}(G) \\ &= 2nr(r-1)^2. \end{aligned}$$

Corollary 10 *Let G be an r-regular connected graph and let α be a whole number. Then,*

1. $M_1^2(L(G)) = \frac{1}{2} \; M_1^2(G)$ *for* $i = 2, j = 0, k = 0$ or $i = 0, j = 2, k = 0$,
2. $M_1^0(L(G)) = \frac{1}{2} \; M_1^0(G) = n$ *for* $i = 0, j = 0, k = 2$,
3. $M_1^1(L(G)) = -2M_1^1(G) = m$ for $i = 1, j = 0, k = 1$ *or* $i = 0, j = 1, k = 1$,
4. $M_1^2(L(G)) = M_1^2(G) = n$ *for* $i = 1, j = 1, k = 0$.

Theorem 18 *Let $K_{r,s}$ be a complete bipartite graph with $1 \leq r \leq s$, and let α be a whole number. Then,*

$$M_1^{\alpha}(L(K_{r,s})) = \frac{1}{2}\sum_{i,j,k\geq 0; i+j+k=\alpha} \frac{\alpha!}{i!\,j!\,k!}\,(-2)^k\; r^{i+1}\times s^{j+1}.$$

Proof Proof is similar to that of previous theorem. □

The following results are immediate from the above theorem.

Corollary 11 *Let $K_{r,s}$ be a complete bipartite graph with $1\leq r\leq s$ and let $\alpha = 2$. Then,*

$$M_1^2(L(K_{r,s})) = rs(r+s-2)^2.$$

Corollary 12 *Let G be an r-regular connected graph and α be a whole number. Then, the sum of all coefficients of $M_1^{i+j+1}(G)$ in the expansion of $M_1^{\alpha}(L(G))$ is equal to zero.*

Corollary 13 *Let G be an r-regular connected graph and α be a whole number. Then, the number of terms in the expansion of $M_1^{\alpha}(L(G))$ is $\binom{\alpha+2}{2}$.*

Let $\delta_1(G)$ be the minimal degree of non-pendant vertices. Now we obtain lower and upper bounds for $M_1^{\alpha}(L(G))$ in terms of the number of pendent vertices and $\delta_1(G)$.

Theorem 19 *For any (n, m)-connected graph G with η pendent vertices and minimal non-pendant vertex degree $\delta_1(G)$,*

$$(\delta_1(G)-1)^{\alpha}[\eta+(m-\eta)2^{\alpha}] \leq M_1^{\alpha}(L(G)) \leq (\Delta(G)-1)^{\alpha}[\eta+(m-\eta)2^{\alpha}].$$

Proof For any $\alpha\geq 0$, we have

$$\begin{aligned} M_1^{\alpha}(L(G)) &= \sum_{uv\in E(G)} [d_G(u)+d_G(v)-2]^{\alpha} \\ &= \sum_{uv\in E(G); d_G(u)=1, d_G(v)\neq 1} [d_G(v)-1]^{\alpha} \\ &\quad + \sum_{uv\in E(G); d_G(u)\neq 1, d_G(v)\neq 1} [d_G(u)+d_G(v)-2]^{\alpha} \\ &\leq \eta(\Delta(G)-1)^{\alpha} + (m-\eta)(2\Delta(G)-2)^{\alpha}. \end{aligned}$$

Thus, the upper bound follows. Similarly,

$$M_1^{\alpha}(L(G)) \geq (\delta_1(G)-1)^{\alpha}[\eta+(m-\eta)2^{\alpha}].$$

Hence, the desired result follows. □

Remark 1 The bounds in the above theorem are attained if and only if $d_G(u) = d_G(v) = \Delta(G) = \delta_1(G)$ for each $uv \in E(G)$ with $d_G(u)\neq 1$, $d_G(v)\neq 1$ and $d_G(v) = \Delta(G) = \delta_1(G)$ for each $uv\in E(G)$ with $d_G(u) = 1$.

4 Jump Graph

The complement of the line graph is called the jump graph $J(G)$ of a graph G, i.e., $J(G) = \overline{L(G)}$. For more details, we refer to [29, 30].

Theorem 20 *For an (n, m)-connected graph G with $n\geq 4$ and $\alpha\geq 0$, we have*

$$M_1^{\alpha}(J(G)) \geq \sum_{i=0}^{\alpha}\binom{\alpha}{i}(-1)^i\,(m-1)^i\,M_1^{\alpha-i}(L(G)).$$

Proof Let G be an (n, m)-connected graph with $n\geq 4$ and $\alpha\geq 0$. Then,

$$\begin{aligned} M_1^{\alpha}(J(G)) &= \sum_{u\in V(J(G))} [d_{J(G)}(u)]^{\alpha} \\ &= \sum_{u\in V(L(G))} \big[(m-1)-d_{L(G)}(u)\big]^{\alpha}, \end{aligned}$$

since $d_{L(G)}(u) + d_{J(G)}(u) = m-1$. Hence,

$$\begin{aligned} M_1^{\alpha}(J(G)) &= \sum_{i=0}^{\alpha}\binom{\alpha}{i}(-1)^i\,(m-1)^i \sum_{u\in V(L(G))} [d_{L(G)}(u)]^{\alpha-i} \\ &= \sum_{i=0}^{\alpha}\binom{\alpha}{i}(-1)^i\,(m-1)^i\,M_1^{\alpha-i}(L(G)). \end{aligned}$$

□

By the above result, we reach to the following results.

Corollary 14 *For any path graph P_n with $n\geq 5$ vertices and $\alpha = 2$, we have*

$$M_1^2(J(P_n)) = n^3 - 9n^2 + 28n - 30.$$

Theorem 21 *Let G be a connected r-regular graph with $\alpha\geq 0$. Then,*

$$\begin{aligned} &M_1^{\alpha}(J(G)) \\ &= nr\sum_{i=0}^{\alpha}\binom{\alpha}{i}(-1)^i\;2^{\alpha-2i-1}\,(r-1)^{\alpha-i}\,(nr-2)^i. \end{aligned}$$

Proof Let G be a connected r-regular graph with $\alpha\geq 0$. Then,

$$\begin{aligned} M_1^{\alpha}(J(G)) &= \sum_{i=0}^{\alpha}\binom{\alpha}{i}(-1)^i\,(m-1)^i\,\frac{nr}{2}\,(2r-2)^{\alpha-i} \\ &= \sum_{i=0}^{\alpha}\binom{\alpha}{i}(-1)^i\left(\frac{nr}{2}-1\right)^i\frac{nr}{2}\,2^{\alpha-i}\,(r-1)^{\alpha-i} \\ &= \sum_{i=0}^{\alpha}\binom{\alpha}{i}(-1)^i\,\frac{(nr-2)^i}{2^i}\,\frac{nr}{2}\,2^{\alpha-i}\,(r-1)^{\alpha-i} \\ &= nr\sum_{i=0}^{\alpha}\binom{\alpha}{i}(-1)^i\;2^{\alpha-2i-1}\,(r-1)^{\alpha-i}\,(nr-2)^i. \end{aligned}$$

□

By the above results, we obtain the followings.

Corollary 15 *For any complete graph K_n with $n \geq 3$ vertices and $\alpha = 2$, we have*

$$M_1^2(J(K_n)) = \frac{n(n-1)}{8}\left[n^4 - 10n^3 + 37n^2 - 60n + 36\right].$$

Corollary 16 *For any cycle C_n with $n \geq 3$ vertices and $\alpha = 2$, we have*

$$M_1^2(J(C_n)) = n^3 - 6n^2 + 9n.$$

Corollary 17 *For any complete bipartite graph $K_{r,s}$ with $2 \leq r \leq s$ and $\alpha = 2$, we have*

$$M_1^2(J(K_{r,s})) = rs\sum_{i=0}^{\alpha}\binom{\alpha}{i}(-1)^i(rs-1)^i(r+s-2)^{\alpha-i}.$$

5 Iterated Line Graph

If G is a graph and $L(G) = L^1(G)$ is its line graph, then $L^k(G)$, $k = 2, 3, \ldots$, defined recursively as $L^k(G) = L(L^{k-1}(G))$, are named as the iterated line graphs of G. For more details, we refer to [31–33].

Theorem 22 *For any complete bipartite graph $K_{r,s}$ with $1 \leq r \leq s$ and $\alpha \geq 0$, we have*

$$M_1^{\alpha}(L^k(K_{r,s})) = \begin{cases} rs(r+s-2)^{\alpha} & \text{for } k = 1; \\ rs\left[\prod_{i=0}^{k-2}\left(2^{i-1}(r+s-2) - 2^i + 1\right)\right] & \\ \quad \times\left(2^{k-1}(r+s-2) - 2^k + 2\right)^{\alpha} & \text{for } k > 1. \end{cases}$$

Proof Let $K_{r,s}$ be a complete bipartite graph with $1 \leq r \leq s$ and $\alpha \geq 0$. By combinatorial methods, we find that $|V(L(K_{r,s}))| = rs$ and $|E(L(K_{r,s}))| = \frac{rs}{2}(r+s-2)$. Since the kth-iterated line graph of $K_{r,s}$ is a $(2^{k-1}(r+s-2) - 2^k + 2)$-regular graph, we also obtain

$$V(L^k(K_{r,s})) = \begin{cases} rs & \text{for } k = 1; \\ rs\left[\prod_{i=0}^{k-2}\left(2^{i-1}(r+s-2) - 2^i + 1\right)\right] & \text{for } k > 1 \end{cases}$$

and

$$E(L^k(K_{r,s})) = \begin{cases} \frac{rs}{2}(r+s-2) & \text{for } k = 1; \\ \frac{rs}{2^k}\left[\prod_{i=0}^{k-1}\left(2^i(r+s-2) - 2^i + 1\right)\right] & \text{for } k > 1 \end{cases}$$

Therefore, $M_1^{\alpha}(L^k(K_{r,s})) = \sum_{u \in V(L^k(K_{r,s}))}\left[d_{L^k(K_{r,s})}(u)\right]^{\alpha}$. Hence, the desired result follows. □

By the above results, we reach to the following.

Corollary 18 *For any star graph $K_{1,s}$ with $s \geq 1$ and $\alpha \geq 2$,*

$$M_1^{\alpha}(L^k(K_{1,s})) = \begin{cases} s(s-1)^{\alpha} & \text{for } k = 1; \\ s\left[\prod_{i=0}^{k-2}\left(2^{i-1}(s-1) - 2^i + 1\right)\right] & \\ \quad \times\left(2^{k-1}(s-1) - 2^k + 2\right)^{\alpha} & \text{for } k > 1. \end{cases}$$

Theorem 23 *Let G be a connected r-regular graph with $\alpha \geq 0$ and $k \geq 1$. Then,*

$$M_1^{\alpha}(L^k(G)) = n\left[\prod_{i=0}^{k-1}\left(2^{i-1}r - 2^i + 1\right)\right]\left(2^k(r-2) + 2\right)^{\alpha}.$$

Proof Let G be a connected r-regular graph with $\alpha \geq 0$ and $k \geq 1$. Then,

$$M_1^{\alpha}(L^k(G)) = \sum_{u \in V(L^k(G))}\left[d_{L^k(G)}(u)\right]^{\alpha}.$$

For a connected r-regular graph G with $|V(G)| = n$, we have

$$|V(L^k(G))| = n\prod_{i=0}^{k-1}\left(2^{i-1}r - 2^i + 1\right),$$

and

$$|E(L^k(G))| = \frac{n}{2^{k+1}}\prod_{i=0}^{k-1}\left(2^i r - 2^{i+1} + 2\right).$$

Further, the kth-iterated line graph of a connected r-regular graph is $\left(2^k r - 2^{k+1} + 2\right)$-regular graph. Therefore,

$$\begin{aligned} M_1^{\alpha}(L^k(G)) &= \sum_{u \in V(L^k(G))}\left[d_{L^k(G)}(u)\right]^{\alpha} \\ &= \sum_{u \in V(L^k(G))}\left[2^k r - 2^{k+1} + 2\right]^{\alpha} \\ &= \sum_{u \in V(L^k(G))}\left[2^k (r-2) + 2\right]^{\alpha} \\ &= n\left[\prod_{i=0}^{k-1}\left(2^{i-1}r - 2^i + 1\right)\right]\left(2^k(r-2) + 2\right)^{\alpha}. \end{aligned}$$

□

By the above results, we finally obtain the following results.

Corollary 19 *For any cycle C_n with $n \geq 3$ and $\alpha \geq 2$, we have*

$$M_1^{\alpha}(L^k(C_n)) = n\,2^{\alpha}.$$

Corollary 20 *For any complete graph K_n with $n \geq 3$ and $\alpha \geq 2$, we have*

$$M_1^{\alpha}(L^k(K_n)) = n\left[\prod_{i=0}^{k-1}\left(2^{i-1}(n-3)+1\right)\right]\left(2^k(n-3)+2\right)^{\alpha}.$$

References

1. Harary F (1969) Graph theory. Addison Wesley, Reading
2. Kulli VR (2012) College graph theory. Vishwa Int Publ, Gulbarga
3. Gutman I, Polansky OE (1986) Mathematical concepts in organic chemistry. Springer, Berlin
4. Todeschini R, Consonni V (2009) Molecular descriptors for chemoinformatics. Wiley, Weinheim
5. Gutman I, Trinajstić N (1972) Graph Theory and molecular orbitals, Total π-electron energy of alternant hydrocarbons. Chem Phys Lett 17:535–538
6. Borovićanin B, Das KC, Furtula B, Gutman I (2017) Zagreb indices: bounds and extremal graphs, bounds in chemical graph theorybasics. Univ. Kragujevac, Kragujevac, pp 67–153
7. Cangul IN, Yurttas A, Togan M, Cevik AS (2017) New formulae for Zagreb indices. In: AIP Conference Proceedings, vol 1863, pp 1–5
8. Das KC, Akgunes N, Togan M, Yurttas A, Cangul IN, Cevik AS (2016) On the first Zagreb index and multiplicative Zagreb coindices of graphs. Analele Stiintifice ale Universitatii Ovidius Constanta 24(1):153–176
9. Gutman I (2013) Degree-based topological indices. Croat Chem Acta 86:351–361
10. Gutman I, Das KC (2014) The first Zagreb index 30 years after. MATCH Commun Math Comput Chem 50:83–92
11. Gutman I, Kulli VR, Chaluvaraju B, Boregowda HS (2017) On Banhatti and Zagreb indices. J Int Math Virtual Inst 7:53–67
12. Miličević A, Nikolić S, Trinajstić N (2004) On reformulated Zagreb indices. Mol Divers 8:393–399
13. Nikolić S, Kovačević G, Miličević A, Trinajstić N (2003) The Zagreb indices 30 years after. Croat Chem Acta 76:113–124
14. Li X, Zhao H (2004) Trees with the first smallest and largest generalized topological indices. MATCH Commun Math Comput Chem 50:57–62
15. Li X, Zheng J (2005) A unified approach to the extremal trees for different indices. MATCH Commun Math Comput Chem 54:195–208
16. Liu B, Gutman I (2007) Estimating the Zagreb and the general Randić indices. MATCH Commun Math Comput Chem 57:617–632
17. Zhang H, Zhang S (2006) Unicyclic graphs with the first three smalllest and largest first general Zagreb index. MATCH Commun Math Comput Chem 55:427–438
18. Furtula B, Gutman I (2015) A forgotten topological index. J Math Chem 53:1184–1190
19. Shirdel GH, Rezapour H, Sayadi AM (2013) The hyper-Zagreb index of graph operations. Iran J Math Chem 4(2):213–220
20. Diaz JB, Metcalf FT (1963) Stronger forms of a class of inequalities of *G*. Bull Am Math Soc 69:415–418
21. Polya G, Szego G (1945) Inequalities for the capacity of a condenser. Am J Math 67:1–32
22. Ruch E, Gutman I (1979) The branching extent of graphs. J Comb Inf Syst 4:285–295
23. Hardy GH, Littlewood JE, Polya G (1952) Inequalities. Cambridge University Press, England
24. Nordhaus EA, Gaddum JW (1956) On complementary graphs. Am Math Mon 63:175–177
25. Aigner M (1969) Graphs whose complement and line graph are isomorphic. J Comb Theory 7:273–275
26. Beineke LW (2006) Derived graphs with derived complement. Recent Trends in Graph Theory, Springer Lecture Notes in Mathematics 186, Berlin, pp 15–24
27. Ranjini PS, Lokesha V, Cangul IN (2011) On the Zagreb indices of the line graphs of the subdivision graphs. Appl Math Comput 218:699–702
28. Koshy T (2003) Discrete mathematics with applications. Academic Press, Cambridge
29. Cvetković DM, Simić SK (1974) Some remarks on the complement of a line graph. Publ Inst Math Beogr 17(31):37–44
30. Hemminger RL, Beineke LW (1978) Line graphs and line digraphs, selected topics in graph theory. Academic Press Inc, Cambridge, pp 271–305
31. Buckley F (1981) Iterated line graphs. Congr Numer 33:390–394
32. Buckley F (1993) The size of iterated line graphs. Graph Theory Notes N Y 25:33–36
33. Ramane HS, Walikar HB, Rao SB, Acharya BD, Hampiholi PR, Jog SR, Gutman I (2005) Spectra and energies of iterated line graphs of regular graphs. Appl Math Lett 18:679–682

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.